arXiv:1108.1288v1 [math.AC] 5 Aug 2011

# Equality of Linear and Symplectic Orbits

Pratyusha Chattopadhyay & Ravi A. Rao
Institute of Mathematical Sciences, C.I.T. Campus, Taramani, Chennai 600 113, India &
Tata Institute of Fundamental Research, Dr. Homi Bhabha Road, Mumbai 400 005, India

*2000 Mathematics Subject Classification: 13C10, 13H05, 15A63, 19A13, 19B10, 19B14*

*Key words: unimodular rows, transvections, elementary symplectic group*

Abstract: It is shown that the set of orbits of the action of the elementary symplectic transvection group on all unimodular elements of a symplectic module over a commutative ring of characteristic not 2 is identical with the set of orbits of the action of the corresponding elementary transvection group. This result is used to get improved injective stability estimates for $K_1$ of the symplectic transvection group over a non-singular affine algebras.

## 1 Introduction

H. Bass introduced two types of linear transvections of a projective module $R \oplus P$ in [1]. He also introduced two types of symplectic transvecitons of a symplectic module $(\mathbb{H}(R) \oplus P, \langle , \rangle \perp \psi_1)$. (These are recalled in §4).

Since elementary automorphisms are homotopic to the identity, we are able to invoke Quillen-Suslin theory (see [10], [13]) to show that

- the groups generated by the two types of elementary linear transformations are the same as the elementary linear group in the free case (see Lemma 4.5).
- the group generated by the two types of elementary symplectic transformations w.r.t. the standard alternating form are the same as the elementary symplectic group in the free case (see Lemma 5.13).

The above generalizes the special case of these results in ([4], Theorem 2).

The title of this paper alludes to the comparison of the elementary linear and elementary symplectic orbits of a unimodular element $(a, b, p)$ in a symplectic module $(\mathbb{H}(R) \perp P)$. In case $P$ is free of rank $\geq 4$, it is established in ([5], Theorem 4.2, Theorem 5.6) that these two orbits coincide. In the Appendix the missing case when $P$ is free of rank 2 is proved by a similar, but slightly more involved argument. This means one has to essentially prove Lemma 2.9 and Lemma 3.1. Note that in the literature (for instance, see [6]. [7]) the general method of the rank 2 symplectic case is usually by recourse to the study of Chevalley groups via the Chevalley commutator formulae; and the approach here is more in the vein of the higher sizes.

Since elementary automorphisms are homotopic to the identity, we show how the Quillen-Suslin machinery enables one to extend the results of [5] to show that the two orbits are equal in the general case when $P$ is a finitely generated projective module. (The transition is by no means automatic!)

## 2 Preliminaries

A row $v = (v_1, \ldots, v_n) \in R^n$ is said to be *unimodular* if there are elements $w_1, \ldots, w_n$ in $R$ such that $v_1w_1 + \cdots + v_nw_n = 1$. $\mathrm{Um}_n(R)$ will denote the set of all unimodular rows $v \in R^n$. Let $I$ be an ideal in $R$. We denote by $\mathrm{Um}_n(R, I)$ the set of all unimodular rows of length $n$ which are congruent to $e_1 = (1, 0, \ldots, 0)$ modulo $I$. (If $I = R$, then $\mathrm{Um}_n(R, I)$ is $\mathrm{Um}_n(R)$).

**Definition 2.1.** Let $P$ be a finitely generated projective $R$-module of rank $n$. An element $v$ in $P$ is said to be *unimodular* if for any maximal ideal $\mathfrak{m}$ of $R$, we have $v_{\mathfrak{m}} \in \mathrm{Um}_n(R_{\mathfrak{m}})$. The collection of unimodular elements of $P$ is denoted by $\mathrm{Um}(P)$. Note that $\mathrm{Um}(P, IP)$ denotes the collection of elements from $\mathrm{Um}(P)$ such that $v_{\mathfrak{m}} \in \mathrm{Um}_n(R_{\mathfrak{m}}, I_{\mathfrak{m}})$. The set $\mathrm{Um}(P, IP)$ is the collection of all *relative unimodular* elements w.r.t. an ideal $I$ of $R$.

The group $\mathrm{GL}_n(R)$ of invertible matrices acts on $R^n$ in a natural way: $v \longrightarrow v\sigma$, if $v \in R^n$, $\sigma \in \mathrm{GL}_n(R)$. This map preserves $\mathrm{Um}_n(R)$, so $\mathrm{GL}_n(R)$ acts on $\mathrm{Um}_n(R)$. Note that any subgroup $\mathrm{G}$ of $\mathrm{GL}_n(R)$ also acts on $\mathrm{Um}_n(R)$. Let $v, w \in \mathrm{Um}_n(R)$, we denote $v \sim_{\mathrm{G}} w$ or $v \in w\mathrm{G}$ if there is a $g \in \mathrm{G}$ such that $v = wg$.

Let $\mathrm{E}_n(R)$ denote the subgroup of $\mathrm{SL}_n(R)$ consisting of all *elementary* matrices, i.e. those matrices which are a finite product of the *elementary generators* $\mathrm{E}_{ij}(\lambda) = I_n + e_{ij}(\lambda)$, $1 \leq i \neq j \leq n$, $\lambda \in R$, where $e_{ij}(\lambda) \in \mathrm{M}_n(R)$ has at most one non-zero entry $\lambda$ in its $(i, j)$-th position.

In the sequel, if $\alpha$ denotes an $m \times n$ matrix, then we let $\alpha^t$ denote its *transpose* matrix. This is of course an $n \times m$ matrix. However, we will mostly be working with square matrices, or rows and columns.

**Definition 2.2. The Relative Groups $\mathbf{E}_n(I)$, $\mathbf{E}_n(R, I)$:** *Let $I$ be an ideal of $R$. The relative elementary group $\mathrm{E}_n(I)$ is the subgroup of $\mathrm{E}_n(R)$ generated as a group by the elements $\mathrm{E}_{ij}(x)$, $x \in I$, $1 \leq i \neq j \leq n$.*

*The relative elementary group $\mathrm{E}_n(R, I)$ is the normal closure of $\mathrm{E}_n(I)$ in $\mathrm{E}_n(R)$.*

*(Equivalently, $\mathrm{E}_n(R, I)$ is generated as a group by $\mathrm{E}_{ij}(a)\mathrm{E}_{ji}(x)E_{ij}(-a)$, with $a \in R$, $x \in I$, $i \neq j$, provided $n \geq 3$* (see [15], Lemma 8)*).*

**Definition 2.3.** $\mathbf{E}_n^1(R, I)$ is the subgroup of $\mathrm{E}_n(R)$ generated by the elements of the form $E_{1i}(a)$ and $E_{i1}(x)$, where $a \in R, x \in I$, and $2 \leq i \leq n$.

**Definition 2.4. Symplectic Group $\mathbf{Sp}_{2n}(R)$:** *This is the isotropy group of the standard symplectic form $\psi_n = \sum\limits_{i=1}^{n} e_{2i-1,2i} - \sum\limits_{i=1}^{n} e_{2i,2i-1}$. In other words, $\mathrm{Sp}_{2n}(R) = \{\alpha \in \mathrm{GL}_{2n}(R) \mid \alpha^t\psi_n\alpha = \psi_n\}$.*

Let $\sigma$ denote the permutation of the natural numbers given by $\sigma(2i) = 2i - 1$ and $\sigma(2i-1) = 2i$.

**Definition 2.5. Elementary Symplectic Group $\mathbf{ESp}_{2n}(R)$:** *We define for $z \in R$, $1 \leq i \neq j \leq 2n$,*

$$se_{ij}(z) = \begin{cases} 1_{2n} + ze_{ij} & \text{if } i = \sigma(j), \\ 1_{2n} + ze_{ij} - (-1)^{i+j} ze_{\sigma(j)\sigma(i)} & \text{if } i \neq \sigma(j) \text{ and } i < j. \end{cases}$$

(It is easy to check that all these generators belong to $\mathrm{Sp}_{2n}(R)$).

*We call them the elementary symplectic matrices over $R$ and the subgroup of $\mathrm{Sp}_{2n}(R)$ generated by them is called the elementary symplectic group $\mathrm{ESp}_{2n}(R)$.*

**Definition 2.6. The Relative Group $\mathbf{ESp}_{2n}(I)$, $\mathbf{ESp}_{2n}(R, I)$:** *Let $I$ be an ideal of $R$. The relative elementary group $\mathrm{ESp}_{2n}(I)$ is the subgroup of $\mathrm{ESp}_{2n}(R)$ generated as a group by the elements $se_{ij}(x)$, $x \in I$ and $1 \leq i \neq j \leq 2n$.*

*The relative elementary group $\mathrm{ESp}_{2n}(R, I)$ is the normal closure of $\mathrm{ESp}_{2n}(I)$ in $\mathrm{ESp}_{2n}(R)$.*

**Definition 2.7.** *The group $\mathrm{ESp}^1_{2n}(R, I)$ is a subgroup of $\mathrm{ESp}_{2n}(R)$ generated by the elements of the form $se_{1i}(a)$ and $se_{j1}(x)$, where $a \in R$, $x \in I$ and $2 \leq i, j \leq 2n$.*

**Notation 2.8.** We fix some notations. $\mathrm{G}(n, R)$ will denote either the linear group $\mathrm{GL}_n(R)$, or the symplectic group $\mathrm{Sp}_{2m}(R)$, **for** $n = 2m$. Also, $\mathrm{E}(n, R)$ will denote either of the elementary subgroups $\mathrm{E}_n(R)$ or $\mathrm{ESp}_{2m}(R)$. The standard elementary generators of $\mathrm{E}(n, R)$ are denoted by $ge_{ij}(a)$, $a \in R$. Let $I$ be an ideal in $R$. Let $\mathrm{G}(n, R, I)$ denote the relative linear groups $\mathrm{GL}_n(R, I)$, or $\mathrm{SL}_n(R, I)$, the relative symplectic group $\mathrm{Sp}_{2m}(R, I)$. Note that $\mathrm{E}(n, I)$ is a subgroup of $\mathrm{E}(n, R)$ generated as a group by the elements $ge_{ij}(x)$, where $x \in I$, and $1 \leq i \neq j \leq n$. Here, $\mathrm{E}(n, R, I)$ denotes the corresponding relative elementary subgroups $\mathrm{E}_n(R, I)$, $\mathrm{ESp}_{2m}(R, I)$, respectively. These are the normal closures of the subgroups $\mathrm{E}(n, I)$ in $\mathrm{E}(n, R)$, which are also known to be generated by the elements $ge_{ij}(a)ge_{ji}(x)ge_{ij}(-a)$, $a \in R, x \in I$, and $1 \leq i \neq j \leq n$ (see [15], Lemma 8). Also, $\mathrm{E}^1(n, R, I)$ is a subgroup of $\mathrm{E}(n, R)$, generated by the elements $ge_{1i}(a)$, where $a \in R$ and $ge_{i1}(x)$, where $x \in I$, $2 \leq i \leq n$ in the linear and symplectic case.

**Lemma 2.9.** *Let $n \geq 3$ in the linear case and $n \geq 4$ in the symplectic case. Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Then the following sequence is exact*

$$1 \longrightarrow \mathrm{E}(n, R, I) \longrightarrow \mathrm{E}^1(n, R, I) \longrightarrow \mathrm{E}^1(n, R/I, 0) \longrightarrow 1.$$

*Thus $\mathrm{E}(n, R, I)$ equals $\mathrm{E}^1(n, R, I) \cap \mathrm{G}(n, R, I)$.*

Proof: For the proof in the linear case reader can see ([8], Lemma 2.2). The proof in the symplectic case is similar when $n \geq 6$. The proof when $n = 4$ is discussed in the Appendix. $\square$

# 3 Local Global Principle for Relative Elementary Group

**Lemma 3.1.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $n \geq 3$ in the linear case and $n \geq 4$ in the symplectic case. Let $\varepsilon = \varepsilon_1 \ldots \varepsilon_r$ be an element in $\mathrm{E}^1(n, R, I)$, where each $\varepsilon_k$ is a (standard) elementary generator. Also $ge_{ij}(Xf(X))$ is a standard elementary generator of $\mathrm{E}^1(n, R[X], I[X])$. Then*

$$\varepsilon \; ge_{ij}(Y^{4^r} X f(Y^{4^r} X)) \; \varepsilon^{-1} \quad = \quad \prod_{t=1}^{s} ge_{i_t j_t}(Y h_t(X, Y)),$$

*where either $i_t = 1$ or $j_t = 1$ and $h_t(X, Y) \in R[X, Y]$, when $i_t = 1$; $h_t(X, Y) \in I[X, Y]$ when $j_t = 1$.*

Proof: Proof in the linear case is straight forward. We establish the result here in the symplectic case when $n \geq 6$. In the Appendix the case when $n = 4$ is also covered.

Given that $ge_{ij}(Xf(X)) \in \mathrm{E}^1(n, R[X], I[X])$. First we assume $i = 1$, hence $f(X) \in R[X]$. We prove the result using induction on $r$, the number of generators of $\varepsilon$. Let $r = 1$ and $\varepsilon = ge_{pq}(a)$. Note that when $p = 1, a \in R$, and when $q = 1, a \in I$. Also note that we use $*$ to represent an element of the ideal $I$.

*Case (1):* Let $(p, q) = (1, j)$. In this case

$$se_{1j}(a) \; se_{1j}(Y^4 X f(Y^4 X)) \; se_{1j}(-a) \quad = \quad se_{1j}(Y^4 X f(Y^4 X)).$$

*Case (2):* Let $(p, q) = (1, \sigma(j))$. In this case

$$\begin{aligned} & se_{1\sigma(j)}(a) \; se_{1j}(Y^4 X f(Y^4 X)) \; se_{1\sigma(j)}(-a) \\ = \quad & se_{12}(* Y^4 X f(Y^4 X)) se_{1j}(Y^4 X f(Y^4 X)). \end{aligned}$$

*Case (3):* Let $(p, q) = (1, k), k \neq j, \sigma(j)$. In this case

$$se_{1k}(a) \; se_{1j}(Y^4 X f(Y^4 X)) \; se_{1k}(-a) \quad = \quad se_{1j}(Y^4 X f(Y^4 X)).$$

*Case (4):* Let $(p, q) = (k, 1), k \neq j, \sigma(j)$. In this case

$$\begin{aligned} & se_{k1}(a) \; se_{1j}(Y^4 X f(Y^4 X)) \; se_{k1}(-a) \\ = \quad & se_{kj}(* Y^4 X f(Y^4 X)) \; se_{1j}(Y^4 X f(Y^4 X)) \\ = \quad & [se_{k1}(* Y^2), \; se_{1j}(Y^2 X f(Y^4 X))] \; se_{1j}(Y^4 X f(Y^4 X)), \end{aligned}$$

where $*$ is an element of $I$.

*Case (5):* Let $(p, q) = (k, 1)$, where either $k = j$ or $k = \sigma(j)$. Let us choose

$r \neq 1, 2, j, \sigma(j)$. In this case

$$\begin{aligned}
& se_{k1}(a)\ se_{1j}(Y^4Xf(Y^4X))\ se_{k1}(-a) \\
= \ & se_{j1}(a)\ [se_{1r}(Y^2Xf(Y^4X)),\ se_{rj}(Y^2)]\ se_{j1}(-a) \\
= \ & [se_{jr}(Y^2X * f(Y^4X))\ se_{1r}(Y^2Xf(Y^4X)),\ se_{r1}(-Y^2*) \\
& se_{rj}(Y^2)] \\
= \ & se_{jr}(Y^2X * f(Y^4X))\ se_{1r}(Y^2Xf(Y^4X))\ se_{r1}(-Y^2*)\ se_{rj}(Y^2) \\
& se_{1r}(-Y^2Xf(Y^4X))\ se_{jr}(-Y^2X * f(Y^4X))\ se_{rj}(-Y^2) \\
& se_{r1}(Y^2*) \\
= \ & se_{jr}(Y^2X * f(Y^4X))\ se_{1r}(Y^2Xf(Y^4X))\ se_{r1}(-Y^2*)\ se_{rj}(Y^2) \\
& se_{1r}(-Y^2Xf(Y^4X))\ [se_{j1}(-Y*), se_{1r}(YXf(Y^4X))]\ se_{rj}(-Y^2) \\
& se_{r1}(Y^2*) \\
= \ & se_{jr}(Y^2X * f(Y^4X))\ se_{1r}(Y^2Yf(Y^4X))\ se_{r1}(-Y^2*)\ se_{rj}(Y^2) \\
& se_{1r}(-Y^2Xf(Y^4X))\ se_{rj}(-Y^2)\ se_{rj}(Y^2) \\
& [se_{j1}(-Y*),\ se_{1r}(YXf(Y^4X))]\ se_{rj}(-Y^2)\ se_{r1}(Y^2*) \\
= \ & [se_{j1}(Y*),\ se_{1r}(YXf(Y^4X))]\ se_{1r}(Y^2Xf(Y^4X))\ se_{r1}(-Y^2*) \\
& se_{1r}(-Y^2Xf(Y^4X))\ se_{1j}(Y^4Xf(Y^4X))\ [se_{r1}(-Y^3*) \\
& se_{j1}(-Y*),\ se_{1j}(-Y^3Xf(Y^4X))\ se_{1r}(YXf(Y^4X))]\ se_{r1}(Y^2*).
\end{aligned}$$

Hence the result is true when $r = 1$. We show the case $j = 1$ by carrying out similar calculations. Let us assume the result is true when the number of generators is $r-1$, i.e, $\varepsilon_2 \ldots \varepsilon_r\ se_{ij}(Y^{4^{r-1}}Xf(Y^{4^{r-1}}(X))\ \varepsilon_r^{-1} \ldots \varepsilon_2^{-1} = \prod_{t=1}^{s} se_{i_tj_t}(Yh_t(X,Y))$, where either $i_t = 1$ or $j_t = 1$. Note that $h_t(X,Y) \in R[X,Y]$, when $i_t = 1$ and $h_t(X,Y) \in I[X,Y]$, when $j_t = 1$.

We prove the result when the number of generators is $r$. We repeat the calculation above to conclude that $\varepsilon se_{ij}(Y^{4^r}Xf(Y^{4^r}X))\varepsilon^{-1} = \prod_{t=1}^{s} se_{p_tq_t}(Yh_t(X,Y))$, where either $p_t = 1$ or $q_t = 1$. Here $h_t(X,Y) \in R[X,Y]$, when $p_t = 1$ and $h_t(X,Y) \in I[X,Y]$, when $q_t = 1$. $\square$

**Lemma 3.2.** *Let $M$ be an $R$-module and $I$ be an ideal of $R$. Let $\alpha(X), \beta(X) \in \mathrm{Aut}(M[X], IM[X]) = ker(\mathrm{Aut}M[X] \longrightarrow \mathrm{Aut}(M[X]/IM[X]))$, with $\alpha(0) = Id$, $\beta(0) = Id$. Let $a$ be a non-nilpotent element in $R$ such that $\alpha(X)_a = \beta(X)_a$ in $\mathrm{Aut}(M_a[X], IM_a[X])$. Then $\alpha(a^NX) = \beta(a^NX)$ in $\mathrm{Aut}(M[X], IM[X])$, for $N \gg 0$.*

Proof: Using $\alpha(0) - \beta(0) = 0$, we get $\alpha(X) - \beta(X) = X\gamma(X)$, for some $\gamma(X)$ in $\mathrm{Aut}(M[X], IM[X])$. Also $\alpha(X)_a - \beta(X)_a = 0$ in $\mathrm{Aut}(M_a[X], IM_a[X])$, i.e, $\big(\alpha(X) - \beta(X)\big)_a = 0$, i.e, $\big(X\gamma(X)\big)_a = 0$. Hence $a^N\big(X\gamma(X)\big) = 0$, in $\mathrm{Aut}(M[X], IM[X])$, for some $N \gg 0$. Therefore $\alpha(a^NX) - \beta(a^NX) = a^NX\gamma(a^NX) = 0$, in $\mathrm{Aut}(M[X], IM[X])$, for $N \gg 0$. $\square$

**Lemma 3.3.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $n \geq 3$ in the linear case and $n \geq 4$ in the symplectic case. Let $a$ be a non-nilpotent element in $R$ and $\alpha(X)$ be in $\mathrm{E}^1(n, R_a[X], I_a[X])$, with $\alpha(0) = Id$. Then there exists $\alpha^*(X) \in \mathrm{E}^1(n, R[X], I[X])$ such that $\alpha^*(X)$ localises to $\alpha(bX)$, for $b \in (a^N)$, where $N \gg 0$, with $\alpha^*(0) = Id$.*

Proof: Let $\alpha(X) = \prod_{k=1}^{r} ge_{i_k j_k}(f_k(X))$, where $f_k(X) = f_k(0) + Xg_k(X)$. Therefore, $\alpha(X) = \prod_{k=1}^{r} \gamma_k \; ge_{i_k j_k}(Xg_k(X)) \; \gamma_k^{-1}$, where $\gamma_l = \prod_{k=1}^{l} ge_{i_k j_k}(f_k(0))$ and $\gamma_l$ is in $\mathrm{E}^1(n, R_a, I_a)$. We have $\alpha(Y^{4^r}X) = \prod_{k=1}^{r} \left( \prod_{t=1}^{s} ge_{i_t j_t}(Yh_t(X,Y))/a^m \right)$, where either $i_t = 1$ or $j_t = 1$, by Lemma 3.1. Note that $h_t(X,Y) \in R[X,Y]$, when $i_t = 1$ and $h_t(X,Y) \in I[X,Y]$, when $j_t = 1$, and $m$ is a natural number. Let us choose $N = m + N'$ and define $\alpha^*(X,Y)$ to be $\prod_{k=1}^{r} \left( \prod_{t=1}^{s} ge_{i_t j_t}(a^{N'} Y h_t(X, a^N Y)) \right)$.

Clearly $\alpha^*(X,Y) \in \mathrm{E}^1(n, R[X,Y], I[X,Y])$ and $\alpha((a^N Y)^{4^r} X) = \alpha^*(X,Y)$. Substituting $Y = 1$, we get $\alpha(bX) = \alpha^*(X)$, for $b \in (a^N)$, $N \gg 0$. Note that $\alpha^*(X) \in \mathrm{E}^1(n, R[X], I[X])$, with $\alpha^*(0) = Id$. $\square$

**Theorem 3.4.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $n \geq 3$ in the linear case and $n \geq 4$ in the symplectic case. Let $a$ be a non-nilpotent element in $R$ and $\alpha(X)$ be in $\mathrm{E}(n, R_a[X], I_a[X])$, with $\alpha(0) = Id$. Then there exists $\alpha^*(X) \in \mathrm{E}(n, R[X], I[X])$ such that $\alpha^*(X)$ localises to $\alpha(bX)$, for $b \in (a^N)$, where $N \gg 0$, with $\alpha^*(0) = Id$.*

Proof: Follows from the previous lemma and Lemma 2.9, which says that $\mathrm{E}(n, R, I)$ is $\mathrm{E}^1(n, R, I) \cap \mathrm{G}(n, R, I)$. $\square$

## 4 Transvection Groups

Following H. Bass one can define transvections of a finitely generated $R$-module as follows:

**Definition 4.1.** Let $M$ be a finitely generated $R$-module. Let $q \in M$ and $\pi \in M^* = \mathrm{Hom}(M, R)$, with $\pi(q) = 0$. Let $\pi_q(p) := \pi(p)q$. An automorphism of the form $1 + \pi_q$ is called a **transvection** of $M$, if either $q \in \mathrm{Um}(M)$ or $\pi \in \mathrm{Um}(M^*)$. Collection of transvections of $M$ is denoted by $\mathrm{Trans}(M)$. This forms a subgroup of $\mathrm{Aut}(M)$.

**Definition 4.2.** Let $M$ be a finitely generated $R$ module. The automorphisms of $N = (R \perp M)$ of the form

$$\begin{aligned} (a,p) &\mapsto (a, p + ax), \\ (a,p) &\mapsto (a + \tau(p), p), \end{aligned}$$

where $x \in M$ and $\tau \in M^*$ are called **elementary transvections** of $N$. Let us denote the first automorphism by $E_x$ and the second one by $E^*_\tau$. It can be verified that these are transvecions of $N$. Let us consider $\pi(t,y) = t$ and $q = (0,x)$ to get $E_x$. Next we can consider $\pi(a,p) = \tau(p)$, where $\tau \in M^*$ and $q = (1,0)$ to get $E^*_\tau$. The subgroup of $\mathrm{Trans}(N)$ generated by elementary transvections is denoted by $\mathrm{ETrans}(N)$.

**Definition 4.3.** Let $I$ be an ideal of $R$. The group of **relative transvections** w.r.t. an ideal $I$ is generated by the transvections of the form $1 + \pi_q$, where either $q \in \mathrm{Um}(IM), \pi \in \mathrm{Um}(M^*)$, or $q \in \mathrm{Um}(M), \pi \in \mathrm{Um}(IM^*)$. The group of relative transvections is denoted by $\mathrm{Trans}(M, IM)$.

**Definition 4.4.** Let $I$ be an ideal of $R$. The elementary transvections of $N = (R \perp M)$ of the form $E_x, E^*_\tau$, where $x \in IM$ and $\tau \in (IM)^*$ are called **relative elementary transvections** w.r.t. an ideal $I$, and the group generated by them is denoted by $\mathrm{ETrans}(IN)$. The normal closure of $\mathrm{ETrans}(IN)$ in $\mathrm{ETrans}(N)$ is denoted by $\mathrm{ETrans}(N, IN)$.

**Lemma 4.5.** *Let $I$ be an ideal of $R$ and $M$ be a free $R$ module of rank $n \geq 2$, and $N = (R \perp M)$. Then* $\mathrm{ETrans}(N, IN) = \mathrm{Trans}(N, IN) = \mathrm{E}_{n+1}(R, I)$.

Proof: Note that when $M$ is a free $R$ module, an element of $\mathrm{Trans}(N, IN)$ looks like $I_{n+1} + v^t w$, for some $v, w \in R^{n+1}$, with either $v$ or $w$ unimodular and belongs to $I^{n+1}$ $(\subseteq R^{n+1})$. Also $\langle v, w\rangle = 0$. Therefore $\mathrm{Trans}(N, IN) \subseteq \mathrm{E}_{n+1}(R, I)$ (follows from [13], Lemma 1.2 and Lemma 1.3).

Using definition of $\mathrm{ETrans}(N, IN)$, $\mathrm{E}_{n+1}(R, I)$ and the fact that in the free case elementary transvections $E_x$ and $E^*_\tau$ of $N$ are of the form $\left(\begin{smallmatrix} 1 & x \\ 0 & I_n \end{smallmatrix}\right)$, and $\left(\begin{smallmatrix} 1 & 0 \\ y^t & I_n \end{smallmatrix}\right)$, respectively, we get $\mathrm{ETrans}(N, IN) \subseteq \mathrm{E}_{n+1}(R, I)$. By ([8], Lemma 2.2) and ([12], Lemma 2.7(a)) we have $\mathrm{E}_{n+1}(R, I) \subseteq \mathrm{ETrans}(N, IN)$, hence $\mathrm{ETrans}(N, IN) = \mathrm{E}_{n+1}(R, I)$. We have $\mathrm{E}_{n+1}(R, I) = \mathrm{ETrans}(N, IN) \subseteq \mathrm{Trans}(N, IN) \subseteq \mathrm{E}_{n+1}(R, I)$, and hence the result follows. $\square$

Here we establish dilation principle for the elementary transvection group.

**Lemma 4.6.** *Let $I$ be an ideal of $R$ and let $M$ be a finitely generated module of $R$. Let $a$ be non-nilpotent element of $R$ such that $M_a$ be free $R_a$-module of rank at least 2. Let $N = (R \perp M)$. Let $\alpha(X) \in \mathrm{ETrans}(N_a[X], IN_a[X])$, with $\alpha(0) = Id$. Then there exists $\alpha^*(X) \in \mathrm{ETrans}(N[X], IN[X])$ such that $\alpha^*(X)$ localises to $\alpha(bX)$, for $b \in (a^N)$, where $N \gg 0$ and $\alpha^*(0) = Id$.*

Proof: Given that $M_a$ is a free $R_a$-module. Using Lemma 4.5 we get that $\mathrm{ETrans}(N_a[X], IN_a[X]) = \mathrm{E}_{n+1}(R_a[X], I_a[X])$. Using Lemma 3.4 we get $\beta^*(X) \in \mathrm{E}_{n+1}(R[X], I[X])$ such that $\beta^*(X)_a = \alpha(b'X)$, where $b' \in (a^N)$, $N \gg 0$.

Let us choose $\alpha^*(X)$ from $\mathrm{ETrans}(N[X], IN[X])$ such that $\alpha^*(X)_a = \beta^*(X)_a$, over the ring $R_a[X]$. Therefore $\alpha^*(X)$ localises to $\alpha(bX)$ for $b \in (a^N)$, where $N \gg 0$ and $\alpha^*(0) = Id$. $\square$

**Lemma 4.7.** *Let $I$ be an ideal of $R$ and let $M$ be a finitely generated projective $R$-module of rank $n \geq 2$. Let $N = (R \perp M)$. Let $\alpha(X) \in \mathrm{Aut}(N[X])$, with $\alpha(0) = Id$. If for each maximal ideal $\mathfrak{m}$ of $R$, $\alpha(X)_\mathfrak{m} \in \mathrm{ETrans}(N_\mathfrak{m}[X], IN_\mathfrak{m}[X])$, then $\alpha(X) \in \mathrm{ETrans}(N[X], IN[X])$.*

Proof: One can suitably choose an element $a_\mathfrak{m}$ from $R \setminus \mathfrak{m}$ such that $\alpha(X)_{a_\mathfrak{m}} \in \mathrm{ETrans}(M_{a_\mathfrak{m}}[X])$. Let us set $\gamma(X, Y) = \alpha(X + Y)_{a_\mathfrak{m}} \alpha(Y)^{-1}_{a_\mathfrak{m}}$. Note that $\gamma(X, Y)$ belongs to $\mathrm{ETrans}(M_{a_\mathfrak{m}}[X, Y], IM_{a_\mathfrak{m}}[X, Y])$, and $\gamma(0, Y) = Id$. From the above lemma it follows that $\gamma(b_\mathfrak{m} X, Y) \in \mathrm{ETrans}(M[X, Y], IM[X, Y])$, for $b_\mathfrak{m} \in (a^N_\mathfrak{m})$, where $N \gg 0$. The ideal generated by $b_\mathfrak{m}$'s is the whole ring $R$. Therefore we have $c_1 b_{\mathfrak{m}_1} + \cdots + c_k b_{\mathfrak{m}_k} = 1$, where $c_i \in R$, for $1 \leq i \leq k$. Note that $\gamma(c_i b_{\mathfrak{m}_i} X, Y)$ belongs to $\mathrm{ETrans}(M[X, Y], IM[X, Y])$, for $1 \leq i \leq k$. Hence $\alpha(X)$ which is equal to $\prod_{i=1}^{k-1} \gamma(c_i b_{\mathfrak{m}_i} X, T_i)\gamma(c_k b_{\mathfrak{m}_k} X, 0)$ is in $\mathrm{ETrans}(M[X], IM[X])$, where $T_i = c_{i+1} b_{\mathfrak{m}_{i+1}} X + \cdots + c_k b_{\mathfrak{m}_k} X$. $\square$

**Theorem 4.8.** *Let $I$ be an ideal of $R$ and let $M$ be a finitely generated projective $R$-module of rank $n \geq 2$. Let $N = (R \perp M)$. Let $q(X) \in \mathrm{Um}(N[X], IN[X])$, where $q(X)$ is of the form $(a(X), p(X))$. If for each maximal ideal $\mathfrak{m}$ of $R$, $q(X) \in q(0)\mathrm{ETrans}(N_{\mathfrak{m}}[X], IN_{\mathfrak{m}}[X])$, then $q(X) \in q(0)\mathrm{ETrans}(N[X], IN[X])$.*

Proof: For each maximal ideal $\mathfrak{m}$ of $R$, we get $\beta_{(\mathfrak{m})}(X) \in \mathrm{ETrans}(M_{\mathfrak{m}}[X], IM_{\mathfrak{m}}[X])$ such that $v(X)\beta_{(\mathfrak{m})}(X) = v(0)$. Let us define $\gamma(X,T) = \beta_{(\mathfrak{m})}(X+T)\beta_{(\mathfrak{m})}(X)^{-1}$. Clearly $\gamma(X,T) \in \mathrm{ETrans}(M_{\mathfrak{m}}[X,T], IM_{\mathfrak{m}}[X,T])$. There are only finitely many denominators involved. Therefore there exists $a_{\mathfrak{m}} \in R \setminus \mathfrak{m}$ such that $\gamma(X,T)$ belongs to $\mathrm{ETrans}(M_{a_{\mathfrak{m}}}[X,T], IM_{a_{\mathfrak{m}}}[X,T])$. Also $\gamma(X,0) = Id$. This implies $\gamma(X, b_{\mathfrak{m}}T) \in \mathrm{ETrans}(M[X,T], IM[X,T])$ for $b_{\mathfrak{m}} \in (a_{\mathfrak{m}}^N)$, $N \gg 0$. Now, $v(X+b_{\mathfrak{m}}T)\gamma(X, b_{\mathfrak{m}}T) = v(X+b_{\mathfrak{m}}T)\beta_{(\mathfrak{m})}(X+b_{\mathfrak{m}}T)\beta_{(\mathfrak{m})}(X)^{-1} = v(0)\beta_{(\mathfrak{m})}(X)^{-1} = v(X)$.

Note that the ideal generated by $b_{\mathfrak{m}}$'s is the whole ring $R$. Therefore $c_1 b_{\mathfrak{m}_1} + \cdots + c_k b_{\mathfrak{m}_k} = 1$, where $c_i \in R$, for $1 \leq i \leq k$. In the above equation replacing $b_{\mathfrak{m}}T$ by $c_1 b_{\mathfrak{m}_1} X$ and $X$ by $c_2 b_{\mathfrak{m}_2} X + \cdots + c_k b_{\mathfrak{m}_k} X$ we get,

$$v(X) = v(b_{\mathfrak{m}_1}X + b_{\mathfrak{m}_2}X + \cdots + b_{\mathfrak{m}_k}X) \in v(b_{\mathfrak{m}_2}X + \cdots + b_{\mathfrak{m}_k}X)\ \mathrm{ETrans}(M[X], IM[X]).$$

Again in the above equation replacing $X$ by $b_{\mathfrak{m}_3}X + \cdots + b_{\mathfrak{m}_k}X$ and $b_{\mathfrak{m}}T$ by $b_{\mathfrak{m}_2}X$ we get, $v(b_{\mathfrak{m}_2}X + \cdots + b_{\mathfrak{m}_k}X) \in v(b_{\mathfrak{m}_3}X + \cdots + b_{\mathfrak{m}_k}X)\mathrm{ETrans}(M[X], IM[X])$. Continuing in this way we get $v(b_{m_k}X + 0) \in v(0)\mathrm{ETrans}(M[X], IM[X])$. Combining all these we get $v(X) \in v(0)\mathrm{ETrans}(M[X], IM[X])$. $\square$

Here we establish equality of two groups.

**Proposition 4.9.** *Let $I$ be an ideal of $R$. Let $M$ be a finitely generated projective $R$-module of rank at least 2, and $N = (R \perp M)$. Then* $\mathrm{Trans}(N, IN) = \mathrm{ETrans}(N, IN)$.

Proof: Note that $\mathrm{ETrans}(N, IN) \subseteq \mathrm{Trans}(N, IN)$. Let us consider an element $\alpha \in \mathrm{Trans}(N, IN)$. There exists $\alpha(X) \in \mathrm{Trans}(N[X], IN[X])$ such that $\alpha(1) = \alpha$ and $\alpha(0) = Id$. Let $\mathfrak{m}$ be a maximal ideal of $R$. We have $\alpha(X)_{\mathfrak{m}} \in \mathrm{Trans}(N_{\mathfrak{m}}[X], IN_{\mathfrak{m}}[X]) = \mathrm{ETrans}(N_{\mathfrak{m}}[X], IN_{\mathfrak{m}}[X])$ (see Lemma 4.5). This is true for all maximal ideal $\mathfrak{m}$ of $R$ and hence by Lemma 4.7 we have $\alpha(X)$ is in $\mathrm{ETrans}(N[X], IN[X])$. Substituting $X = 1$ we get $\alpha \in \mathrm{ETrans}(N, IN)$, and hence $\mathrm{Trans}(N, IN) \subseteq \mathrm{ETrans}(N, IN)$. $\square$

## 5 Symplectic Modules and Symplectic Transvections

**Definition 5.1. Alternating Matrix:** *A matrix in $\mathrm{M}_n(R)$ is said to be alternating if it has the form $\nu - \nu^t$, where $\nu \in \mathrm{M}_n(R)$.* (It follows that its diagonal elements are zeros.)

**Lemma 5.2.** *Let $(R, \mathfrak{m})$ be a local ring and $I$ be an ideal of $R$. Let $\varphi$ be an alternating matrix of Pfaffian 1 over $R$, and $\varphi \equiv \psi_n \pmod{I}$. Then $\varphi$ is of the form*

$$(1 \perp \varepsilon)^t \psi_n (1 \perp \varepsilon),$$

*for some $\varepsilon \in \mathrm{E}_{2n-1}(R, I)$.*

Proof: Follows using induction on n and the fact that for any $v \in \mathrm{Um}_n(R, I)$, $v = e_1\beta$, for some $\beta \in \mathrm{E}_n(R, I)$. $\square$

**Remark 5.3.** *The condition that the alternating matrices, in this article, are of Pfaffian one can be extended to all invertible alternating matrices by observing that an invertible alternating matrix over a local ring which is congruent to* $(u\,\psi_1 \perp \psi_{n-1})$ $(\mathrm{mod}\ I)$*, where* $u =$ *Pfaffian* $\varphi$*, is of the form* $(1 \perp E)^t(u\psi_1 \perp \psi_{n-1})(1 \perp E)$*, for some relative elementary matrix* $E$*. Only slight modifications in the proofs given below are needed, which is an easy exercise.*

**Remark 5.4.** *Let* $\varphi$ *be an alternating matrix of Pfaffian* $1$*, over* $R$*. Let us consider the local ring* $R_{\mathfrak{m}}$*, where* $\mathfrak{m}$ *be a maximal ideal of* $R$*. We will get* $\varepsilon(\mathfrak{m}) \in \mathrm{E}_{2n-1}(R_{\mathfrak{m}})$ *such that over* $R_{\mathfrak{m}}$ *we have* $\varphi = (1 \perp \varepsilon(\mathfrak{m}))^t\psi_n(1 \perp \varepsilon(\mathfrak{m}))$*. Let* $a$ *be the product of denominators of all the entries of* $\varepsilon(\mathfrak{m})$*. Clearly* $a$ *is not in* $\mathfrak{m}$*. Hence we get* $\varepsilon$ *from* $\mathrm{E}_{2n-1}(R_a)$ *such that* $\varphi = (1 \perp \varepsilon)^t\psi_n(1 \perp \varepsilon)$*.*

*Note: When dealing with relative case w.r.t. an ideal* $I$ *of* $R$*, we will always assume that the alternating matrix* $\varphi$ *of Pfaffian 1 is congruent to* $\psi_n(\mathrm{mod}\ I)$*. Using Lemma 5.2 and arguing as above we get that over ring* $R_{\mathfrak{m}}$ *we have* $\varphi = (1 \perp \varepsilon)^t\psi_n(1 \perp \varepsilon)$*, where* $\varepsilon \in \mathrm{E}_{2n-1}(R_a, I_a)$*, for some* $a \notin \mathfrak{m}$*. We will constantly use this fact without even refering to it!*

**Definition 5.5.** A **symplectic** $R$**-module** is a pair $(P, \langle , \rangle)$, where $P$ is a finitely generated projective $R$-module of even rank and $\langle , \rangle : P \times P \longrightarrow R$ is a non-degenerate (i.e, $P \cong P^*$ by $x \longrightarrow \langle x, \rangle$) alternating bilinear form.

**Definition 5.6.** Let $(P_1, \langle , \rangle_1)$ and $(P_2, \langle , \rangle_2)$ be two symplectic $R$-modules. Their **orthogonal sum** is the pair $(P, \langle , \rangle)$, where $P = P_1 \oplus P_2$ and the inner product is defined by $\langle (p_1, p_2), (q_1, q_2) \rangle = \langle p_1, q_1 \rangle_1 + \langle p_2, q_2 \rangle_2$.

There is a unique non-degenerate bilinear form $\langle , \rangle$ on the $R$-module $\mathbb{H}(R) = R \oplus R^*$, namely $\langle (a_1, b_1), (a_2, b_2) \rangle = a_1b_2 - a_2b_1$.

Now onwards $Q$ will denote $(R^2 \perp P)$ with induced form on $(\mathbb{H}(R) \perp P)$, and $Q[X]$ will denote $(R[X]^2 \perp P[X])$ with induced form on $(\mathbb{H}(R[X]) \perp P[X])$.

**Definition 5.7.** An **isometry** of a symplectic module $(P, \langle , \rangle)$ is an automorphism of $P$ which fixes the bilinear form. The group of isometries of $(P, \langle , \rangle)$ is denoted by $\mathrm{Sp}(P, \langle , \rangle)$.

**Definition 5.8.** In [3] Bass has defined a **symplectic transvection** of a symplectic module $P$ to be an automorphism of the form

$$\sigma(p) \quad = \quad p + \langle u, p \rangle v + \langle v, p \rangle u + \alpha \langle u, p \rangle u,$$

where $\alpha \in R$ and $u, v \in P$ are fixed elements with $\langle u, v \rangle = 0$. It is easy to check that $\langle \sigma(p), \sigma(q) \rangle = \langle p, q \rangle$ and $\sigma$ has an inverse $\tau(p) = p - \langle u, p \rangle v - \langle v, p \rangle u - \alpha \langle u, p \rangle u$.

The subgroup of $\mathrm{Sp}(P, \langle , \rangle)$ generated by the symplectic transvections is denoted by $\mathrm{Trans}_{\mathrm{Sp}}(P, \langle , \rangle)$ (see [14], Page 35).

**Definition 5.9.** The symplectic transvections of $Q = (R^2 \perp P)$ of the form

$$\begin{aligned}(a,b,p) &\mapsto (a, b - \langle p,q\rangle + \alpha a, p + aq),\\ (a,b,p) &\mapsto (a + \langle p,q\rangle - \beta b, b, p + bq),\end{aligned}$$

where $\alpha, \beta \in R$ and $q \in P$, are called **elementary symplectic transvections**.

The elementary symplectic transvections are symplectic transvections on $Q$. Take $(u,v) = ((0,1,0),(0,0,q))$ and $(u,v) = ((-1,0,0),(0,0,q))$ respectively to get the above two transvections of $Q$.

The subgroup of $\mathrm{Trans}_{\mathrm{Sp}}(Q, \langle,\rangle)$ generated by elementary symplectic transvections is denoted by $\mathrm{ETrans}_{\mathrm{Sp}}(Q, \langle,\rangle)$.

**Definition 5.10.** Let $I$ be an ideal of $R$. The group of **relative symplectic transvections** w.r.t. an ideal $I$ is generated by the symplectic transvecions of the form $\sigma(p) = p + \langle u,p\rangle v + \langle v,p\rangle u + \alpha\langle u,p\rangle u$, where $\alpha \in I$ and $u \in P$, $v \in IP$ are fixed elements with $\langle u,v\rangle = 0$. The group generated by relative symplectic transvections is denoted by $\mathrm{Trans}_{\mathrm{Sp}}(P, IP, \langle,\rangle)$.

**Definition 5.11.** The elementary symplectic transvections of $Q$ of the form

$$\begin{aligned}(a,b,p) &\mapsto (a, b - \langle p,q\rangle + \alpha a, p + aq),\\ (a,b,p) &\mapsto (a + \langle p,q\rangle - \beta b, b, p + bq),\end{aligned}$$

where $\alpha, \beta \in I$ and $q \in IP$ are called **relative elementary symplectic transvections** w.r.t. an ideal $I$.

The subgroup of $\mathrm{ETrans}_{\mathrm{Sp}}(Q, \langle,\rangle)$ generated by relative elementary symplectic transvections is denoted by $\mathrm{ETrans}_{\mathrm{Sp}}(IQ, \langle,\rangle)$. The normal closure of $\mathrm{ETrans}_{\mathrm{Sp}}(IQ, \langle,\rangle)$ in $\mathrm{ETrans}_{\mathrm{Sp}}(Q, \langle,\rangle)$ is denoted by $\mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle,\rangle)$ .

**Remark 5.12.** *Let $P$ be a free $R$-module and $\langle p,q\rangle = p\varphi q^t$, where $\varphi$ be an alternating matrix with Pfaffian 1.*

*In this case the symplectic transvection $\sigma(p) = p + \langle u,p\rangle v + \langle v,p\rangle u + \alpha\langle u,p\rangle u$ corresponds to the matrix $(I_{2n} - v^t u\varphi - u^t v\varphi)(I_{2n} - \alpha u^t u\varphi)$ and the group generated by them is denoted by* $\mathrm{Trans}_{\mathrm{Sp}}(P, \langle,\rangle_\varphi)$.

*Also in this case* $\mathrm{ETrans}_{\mathrm{Sp}}(Q, \langle,\rangle_\varphi)$ *will be generated by the matrices of the form* $\rho_\varphi(q,\alpha) = \begin{pmatrix} 1 & 0 & 0 \\ \alpha & 1 & q\varphi \\ q^t & 0 & I_{2n} \end{pmatrix}$, *and* $\mu_\varphi(q,\beta) = \begin{pmatrix} 1 & -\beta & -q\varphi \\ 0 & 1 & 0 \\ 0 & q^t & I_{2n} \end{pmatrix}$.

*Note that for $q = (q_1, \ldots, q_{2n}) \in R^{2n}$, and for the standard alternating matrix $\psi_n$, we have*

$$\rho_\psi(q,\alpha) = se_{21}(\alpha)\prod_{i=3}^{2n+2} se_{i1}(q_{i-2}), \tag{1}$$

$$\mu_\psi(q,\beta) = se_{12}(\beta)\prod_{i=3}^{2n+2} se_{1i}((-1)^i q_{\sigma(i-2)}). \tag{2}$$

**Lemma 5.13.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $P$ be a free $R$-module of rank $2n$, $n \geq 1$. Let $\varphi = \psi_n$, the standard alternating matrix, then* $\mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_n}) = \mathrm{ESp}_{2n+2}(R, I)$.

Proof: We first show $\mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_n})$ is a subset of $\mathrm{ESp}_{2n+2}(R, I)$. It is enough to show that an element of the form $t_{\psi_n}(q_1, \alpha)s_{\psi_n}(q_2, \beta)t_{\psi_n}(q_1, \alpha)^{-1}$ is in $\mathrm{ESp}_{2n}(R, I)$, for $q_1 \in R^{2n}$, $q_2 \in I^{2n}$, $\alpha \in R$ and $\beta \in I$. Here $t_{\psi_n}$ and $s_{\psi_n}$ are $\rho_{\psi_n}$ or $\mu_{\psi_n}$. Using equations (1), (2) we get $\mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_n}) \subseteq \mathrm{ESp}_{2n}(R, I)$.

To show the other inclusion we recall the equivalent definition of the relative group which says that $\mathrm{ESp}_{2n}(R, I)$ is the smallest normal subgroup of $\mathrm{ESp}_{2n}(R)$ containing $se_{21}(x)$, where $x \in I$ (see [8], Lemma 2.2). We have $gse_{21}(x)g^{-1} \in \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_n})$, hence $\mathrm{ESp}_{2n}(R, I) \subseteq \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_n})$, and hence the equality is established. □

**Lemma 5.14.** *Let $P$ be a free $R$-module of rank $2n$. Let $(P, \langle , \rangle_\varphi)$ and $(P, \langle , \rangle_{\varphi^*})$ be two symplectic $R$-modules with $\varphi = (1 \perp \varepsilon)^t \ \varphi^* \ (1 \perp \varepsilon)$, for some $\varepsilon \in \mathrm{E}_{2n-1}(R)$. Then*

$$\begin{aligned} \mathrm{Trans}_{\mathrm{Sp}}(P, \langle , \rangle_\varphi) &= (1 \perp \varepsilon)^{-1} \ \mathrm{Trans}_{\mathrm{Sp}}(P, \langle , \rangle_{\varphi^*}) \ (1 \perp \varepsilon), \\ \mathrm{ETrans}_{\mathrm{Sp}}(Q, \langle , \rangle_\varphi) &= (I_3 \perp \varepsilon)^{-1} \ \mathrm{ETrans}_{\mathrm{Sp}}(Q, \langle , \rangle_{\varphi^*}) \ (I_3 \perp \varepsilon)). \end{aligned}$$

Proof: In the free case for symplectic transvections we have

$$\begin{aligned} & (I_{2n} - v^t u\varphi - u^t v\varphi)(I_{2n} - \alpha u^t u\varphi) \\ = \ & (1 \perp \varepsilon)^{-1} \ (I_{2n} - \tilde{v}^t \tilde{u}\varphi^* - \tilde{u}^t \tilde{v}\varphi^*) \ (I_{2n} - \alpha \tilde{u}^t \tilde{u}\varphi^*) \ (1 \perp \varepsilon), \end{aligned}$$

where $\tilde{u} = u(1 \perp \varepsilon)^t$ and $\tilde{v} = v(1 \perp \varepsilon)^t$. Hence the first equality follows.

For elementary symplectic transvections we have

$$\begin{aligned} & (I_2 \perp (1 \perp \varepsilon))^{-1} \rho_{\varphi^*}(q, \alpha)(I_2 \perp (1 \perp \varepsilon)) \\ = \ & \rho_\varphi(q(1 \perp \varepsilon^t)^{-1}, \alpha), \\ & (I_2 \perp (1 \perp \varepsilon))^{-1} \mu_{\varphi^*}(q, \beta)(I_2 \perp (1 \perp \varepsilon)) \\ = \ & \mu_\varphi(q(1 \perp \varepsilon^t)^{-1}, \beta), \end{aligned}$$

hence the second equality follows. □

**Lemma 5.15.** *Let $I$ be an ideal of $R$ and $P$ be a free $R$-module of rank $2n$. Let $(P, \langle , \rangle_\varphi)$ and $(P, \langle , \rangle_{\varphi^*})$ be two symplectic $R$-modules with $\varphi = (1 \perp \varepsilon)^t \ \varphi^* \ (1 \perp \varepsilon)$, for some $\varepsilon \in \mathrm{E}_{2n-1}(R, I)$. Then*

$$\begin{aligned} \mathrm{Trans}_{\mathrm{Sp}}(P, IP, \langle , \rangle_\varphi) &= (1 \perp \varepsilon)^{-1} \ \mathrm{Trans}_{\mathrm{Sp}}(P, IP, \langle , \rangle_{\varphi^*}) \ (1 \perp \varepsilon), \\ \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_\varphi) &= (I_3 \perp \varepsilon)^{-1} \ \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\varphi^*}) \ (I_3 \perp \varepsilon). \end{aligned}$$

Proof: Using the three equations appear in the proof of Lemma 5.14, we get these equalities. □

**Proposition 5.16.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $(P, \langle , \rangle_{\varphi})$ be a symplectic $R$-module with $P$ free of rank $2n$, $n \geq 1$. Let $\varphi = (1 \perp \varepsilon)^t\ \psi_n\ (1 \perp \varepsilon)$, for some $\varepsilon \in \mathrm{E}_{2n-1}(R, I)$. Then* $\mathrm{Trans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_1 \perp \varphi}) = \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\varphi})$.

Proof: Using Lemma 5.13 and Lemma 5.15 we get,

$$\begin{aligned} \mathrm{Trans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_1 \perp \varphi}) &= (I_3 \perp \varepsilon)^{-1}\ \mathrm{Trans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_1 \perp \psi_n})\ (I_3 \perp \varepsilon) \\ &= (I_3 \perp \varepsilon)^{-1}\ \mathrm{ESp}_{2+2n}(R, I)\ (I_3 \perp \varepsilon), \end{aligned}$$

and

$$\begin{aligned} \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\varphi}) &= (I_3 \perp \varepsilon)^{-1}\ \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_n})\ (I_3 \perp \varepsilon) \\ &= (I_3 \perp \varepsilon)^{-1}\ \mathrm{ESp}_{2+2n}(R, I)\ (I_3 \perp \varepsilon), \end{aligned}$$

and hence the sequence of equalities are established. □

**Remark 5.17.** *In view of above two lemmas, for any symplectic module $(P, \langle , \rangle_{\varphi})$ over a local ring $(R, \mathfrak{m})$, we have* $\mathrm{Trans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\psi_1 \perp \varphi}) = \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle_{\varphi})$. *Here $I$ is an ideal of the ring $R$.*

Next we establish dilation principle for elementary symplectic transvection group.

**Lemma 5.18.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $(P, \langle , \rangle)$ be a symplectic $R$-module with $P$ finitely generated projective $R$-module of rank $2n$, $n \geq 1$. Let $a \in R$ be non-nilpotent and $(P_a, \langle , \rangle_{\varphi})$ be a symplectic module with $P_a$ be a free $R_a$-module. Also let $\varphi = (1 \perp \varepsilon)^t\ \psi_n\ (1 \perp \varepsilon)$, for some $\varepsilon \in \mathrm{E}_{2n-1}(R_a, I_a)$ over the ring $R_a$. Let $\alpha(X) \in \mathrm{ETrans}_{\mathrm{Sp}}(Q_a[X], IQ_a[X], \langle , \rangle_{\varphi})$, with $\alpha(0) = Id$. Then there exists $\alpha^*(X) \in \mathrm{ETrans}_{\mathrm{Sp}}(Q[X], IQ[X], \langle , \rangle)$ such that $\alpha^*(X)$ localises to $\alpha(bX)$, where $b \in (a^N)$, for $N \gg 0$, and $\alpha^*(0) = Id$.*

Proof: We have

$$\begin{aligned} & \mathrm{ETrans}_{\mathrm{Sp}}(Q_a[X], IQ_a[X], \langle , \rangle_{\varphi}) \\ = \ & (I_3 \perp \varepsilon)^{-1} \mathrm{ETrans}_{\mathrm{Sp}}(Q_a[X], IQ_a[X], \langle , \rangle_{\psi_n})(I_3 \perp \varepsilon) \\ = \ & (I_3 \perp \varepsilon)^{-1} \mathrm{ESp}_{2n+2}(R_a[X], I_a[X])(I_3 \perp \varepsilon). \end{aligned}$$

Therefore $\alpha(X) = (I_3 \perp \varepsilon)^{-1}\beta(X)(I_3 \perp \varepsilon)$, where $\beta(X) \in \mathrm{ESp}_{2n+2}(R_a[X], I_a[X])$. Also, $\beta(0) = Id$. Note that $\mathrm{ESp}_{2n+2}(R_a[X], I_a[X]) = \mathrm{ESp}^1_{2n+2}(R_a[X], I_a[X]) \cap \mathrm{Sp}_{2n+2}(R_a[X], I_a[X])$. Using Lemma 3.1 we get $\beta(XY^{4^r}) = \prod se_{i_k j_k}(Yh_k(X,Y)/a^m)$, for $r \gg 0$ and for each $se_{i_k j_k}$ either $i_k$ is 1 or $j_k$ is 1. Hence $\beta(XY^{4^r})$ is equal to $\prod_k \tau_{\psi_n}\big(Yq_{k1}(X,Y)/a^m, Yg_{k2}(X,Y)/a^m\big)$, where $\tau_{\psi_n}$ is either $\rho_{\psi_n}$ or $\mu_{\psi_n}$. Therefore $\alpha(XY^{4^r}) = \prod_k \tau_{\varphi}\big((Yq_{k1}(X,Y)/a^m)(1 \perp \varepsilon^t)^{-1}, (Yg_{k2}(X,Y)/a^m)(1 \perp \varepsilon^t)^{-1}\big)$, where $\tau_{\varphi}$ is either $\rho_{\varphi}$ or $\mu_{\varphi}$ (see Lemma 5.14). Let us choose $\alpha^*(X, Y)$ from the group $\mathrm{ETrans}_{\mathrm{Sp}}(Q[X,Y], IQ[X,Y], \langle , \rangle_{\varphi})$ such that $\alpha^*(X,Y)_a = \alpha(X(a^N Y^{4^r}))$, for $N \gg 0$. Substituting $Y = 1$ we get $\alpha^*(X)_a = \alpha(a^N X)$ □

**Lemma 5.19.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $(P, \langle,\rangle)$ be a symplectic $R$-module with $P$ finitely generated projective module of rank $2n$, $n \geq 1$. Let $\alpha(X) \in \mathrm{Sp}(Q[X], \langle.\rangle)$, with $\alpha(0) = Id$. If for each maximal ideal $\mathfrak{m}$ of $R$, $\alpha(X)_{\mathfrak{m}} \in \mathrm{ETrans}_{\mathrm{Sp}}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X], \langle,\rangle_{\varphi_{\mathfrak{m}}})$, where $\varphi_{\mathfrak{m}} \equiv \psi_n \pmod{I}$, then $\alpha(X) \in \mathrm{ETrans}_{\mathrm{Sp}}(Q[X], IQ[X], \langle,\rangle)$.*

Proof: Proof follows arguing similarly as in Lemma 4.7. ☐

**Theorem 5.20.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $(P, \langle,\rangle)$ be a symplectic $R$-module with $P$ finitely generated projective module of rank $2n$, $n \geq 1$. Let $q(X) = (a(X), b(X), p(X))$ is in $\mathrm{Um}(Q[X], IQ[X])$. If for each maximal ideal $\mathfrak{m}$ of $R$, we have $q(X) \in q(0)\mathrm{ETrans}_{\mathrm{Sp}}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X], \langle,\rangle_{\varphi_{\mathfrak{m}}})$, where $\varphi_{\mathfrak{m}} \equiv \psi_n \pmod{I}$, then $q(X) \in q(0)\mathrm{ETrans}_{\mathrm{Sp}}(Q[X], IQ[X], \langle,\rangle)$.*

Proof: Proof follows arguing similarly as in Theorem 4.8. ☐

Here we establish equality of the symplectic transvection group and the elementary symplectic transvection group.

**Theorem 5.21.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $(P, \langle,\rangle)$ be a symplectic $R$-module with $P$ finitely generated projective module of rank $2n$, $n \geq 1$. Also assume for any maximal ideal $\mathfrak{m}$ of $R$, the alternating form $\langle,\rangle$ corresponds to the alternating matrix $\varphi_{\mathfrak{m}}$, where $\varphi_{\mathfrak{m}} \equiv \psi_n \pmod{I}$, over the ring $R_{\mathfrak{m}}$. Then $\mathrm{Trans}_{\mathrm{Sp}}(Q, IQ, \langle,\rangle) = \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle,\rangle)$.*

Proof: We have $\mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle,\rangle) \subseteq \mathrm{Trans}_{\mathrm{Sp}}(Q, IQ, \langle,\rangle)$. We need to show other inclusion. Let us choose $\alpha$ from $\mathrm{Trans}_{\mathrm{Sp}}(Q, IQ, \langle,\rangle)$. There exists $\alpha(X)$ in $\mathrm{Trans}_{\mathrm{Sp}}(Q[X], IQ[X], \langle,\rangle_{\varphi \otimes R[X]})$ such that $\alpha(1) = \alpha$ and $\alpha(0) = Id$. For each maximal ideal, $\mathrm{Trans}_{\mathrm{Sp}}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X], \langle,\rangle_{\varphi_{\mathfrak{m}}}) = \mathrm{ETrans}_{\mathrm{Sp}}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X], \langle,\rangle_{\varphi_{\mathfrak{m}}})$ (see Remark 5.17). Hence $\alpha(X)_{\mathfrak{m}} \in \mathrm{ETrans}_{\mathrm{Sp}}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X], \langle,\rangle_{\varphi \otimes R_{\mathfrak{m}}[X]})$, for each maximal ideal $\mathfrak{m}$ of $R$. Therefore from Lemma 5.19 it follows that $\alpha(X)$ is in $\mathrm{ETrans}_{\mathrm{Sp}}(Q[X], IQ[X], \langle,\rangle)$ and hence substituting $X = 1$ we get the result. ☐

## 6 Equality of orbits

In this section we establish main result of this article regarding equality of orbits.

**Theorem 6.1.** *Let $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Let $(P, \langle,\rangle)$ be a symplectic $R$-module with $P$ finitely generated projective module of rank $2n$, $n \geq 1$. Let $v = (a, b, p) \in \mathrm{Um}(Q, IQ)$. Then*

$$(a, b, p)\ \mathrm{ETrans}(Q, IQ) \quad = \quad (a, b, p)\ \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle,\rangle).$$

*Here we also assume that for each maximal ideal $\mathfrak{m}$ of $R$, the alternating form $\langle,\rangle$ corresponds to the alternating matrix $\varphi_{\mathfrak{m}}$, where $\varphi_{\mathfrak{m}} \equiv \psi_n \pmod{I}$, over the local ring $R_{\mathfrak{m}}$.*

Proof: Let $\alpha \in \mathrm{ETrans}(Q, IQ)$. Let us choose $\alpha(X)$ from $\mathrm{ETrans}(Q[X], IQ[X])$ such that $\alpha(1) = \alpha$ and $\alpha(0) = Id$. Let us define $V(X) = (a, b, p)\alpha(X)$.

Let $\mathfrak{m}$ be a maximal ideal of $R$. Over $R_{\mathfrak{m}}$, we have $\varphi_{\mathfrak{m}} = (1 \perp \varepsilon(\mathfrak{m}))^t\ \psi_n\ (1 \perp \varepsilon(\mathfrak{m}))$, where $\varepsilon(\mathfrak{m}) \in \mathrm{E}_{2n}(R_{\mathfrak{m}}, I_{\mathfrak{m}})$. Let us define $W(X) = V(X)\ (1 \perp \varepsilon(\mathfrak{m}))^{-1}$. We have

$$\begin{aligned} W(X) &\in W(0)\ \mathrm{E}_{2n+2}(R_{\mathfrak{m}}[X], I_{\mathfrak{m}}[X]) \\ &= W(0)\ \mathrm{ESp}_{2n+2}(R_{\mathfrak{m}}[X], I_{\mathfrak{m}}[X]) \\ &= W(0)\ \mathrm{ETrans}_{\mathrm{Sp}}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X]), \langle , \rangle_{\psi_n}), \end{aligned}$$

and hence $V(X) \in V(0)\ \mathrm{ETrans}_{\mathrm{Sp}}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X], \langle , \rangle_{\varphi_{\mathfrak{m}}})$. The first equality above follows from ([5], Theorem 5.6) when $n > 2$. For $n = 2$ the result has been proved in the Appendix — see Theorem 7.8. This is true for all maximal ideal $\mathfrak{m}$ of $R$, hence by Theorem 5.20, $V(X) \in V(0)\ \mathrm{ETrans}_{\mathrm{Sp}}(Q[X], I[X], \langle , \rangle)$. Substituting $X = 1$ we get

$$(a, b, p)\ \alpha \quad \in \quad (a, b, p)\ \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle).$$

Now we show the other inclusion. Let us consider $\beta$ from $\mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ, \langle , \rangle)$. Let $\beta(X)$ be an element of $\mathrm{ETrans}_{\mathrm{Sp}}(Q[X], IQ[X], \langle , \rangle)$ such that $\beta(1) = \beta$ and $\beta(0) = Id$. We define $V(X) = (a, b, p)\beta(X)$.

Let $\mathfrak{m}$ be a maximal ideal of $R$. Over the local ring $R_{\mathfrak{m}}$, we define $W(X) = V(X)\ (1 \perp \varepsilon(\mathfrak{m}))^{-1}$. We have

$$\begin{aligned} W(X) &\in W(0)\ (1 \perp \varepsilon(\mathfrak{m}))\ \mathrm{ETrans}_{\mathrm{Sp}}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X], \langle , \rangle_{\varphi_{\mathfrak{m}}})\ (1 \perp \varepsilon(\mathfrak{m}))^{-1} \\ &= W(0)\ \mathrm{ETrans}_{\mathrm{Sp}}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X], \langle , \rangle_{\psi_n}) \\ &= W(0)\ \mathrm{ESp}_{2n+2}(R_{\mathfrak{m}}[X], I_{\mathfrak{m}}[X]) \\ &= W(0)\ \mathrm{E}_{2n+2}(R_{\mathfrak{m}}[X], I_{\mathfrak{m}}[X]), \end{aligned}$$

and hence $V(X) \in V(0)\ \mathrm{ETrans}(Q_{\mathfrak{m}}[X], IQ_{\mathfrak{m}}[X])$. Here the last equality follows from Theorem 7.8. This is true for all maximal ideal $\mathfrak{m}$ of $R$, hence by Theorem 4.8, $V(X) \in V(0)\mathrm{ETrans}(Q[X], IQ[X])$. Substituting $X = 1$, we get

$$(a, b, p)\ \beta \quad \in \quad (a, b, p)\ \mathrm{ETrans}(Q, IQ).$$

$\square$

# 7 Decrease in injective stability for $\mathrm{Sp}(Q, \langle , \rangle)/\mathrm{ETrans}_{\mathrm{Sp}}(Q, \langle , \rangle)$

Final goal of this section is to give an improvement for Basu-Rao ([4], Theorem 2) estimate in the module case over finitely generated rings. For this purpose we state and prove a few preliminary results. While dealing with the results in the relative case w.r.t. an ideal $I$ of a ring $R$, we will always assume that over the local ring $R_{\mathfrak{m}}$, where $\mathfrak{m}$ is a maximal ideal, the alternating form $\langle , \rangle$ corresponds to the alternating matrix $\varphi_{\mathfrak{m}}$, where $\varphi_{\mathfrak{m}} \equiv \psi_n \pmod{I}$.

**Theorem 7.1.** ([2], Theorem 3.4, Page 183) *Let $R$ be a commutative ring of dim $d$. Let $I$ be an ideal of $R$ and $P$ be a projective module of rank $\geq d+1$. Let $\tilde{Q} = R \perp P$. Let $v_1, v_2 \in \mathrm{Um}(\tilde{Q})$ and $v_1 \equiv v_2 \pmod{I\tilde{Q}}$. Then there exists $\beta \in \mathrm{ETrans}(\tilde{Q}, I\tilde{Q})$ such that $v_1\beta = v_2$.* $\square$

**Lemma 7.2.** *Let $R$ be commutative ring of dimension $d$, and let $I$ be an ideal of $R$. Let us assume $R = 2R$. Let $(P, \langle , \rangle)$ be a symplectic $R$-module with $P$ finitely generated projective module of even rank $\geq max\{2, d\}$, and let $Q = R^2 \perp P$. Let $v_1, v_2 \in \mathrm{Um}(Q)$ and $v_1 \equiv v_2 \pmod{IQ}$. Then there exists $\beta \in \mathrm{ETrans}_{\mathrm{Sp}}(Q, IQ)$ such that $v_1\beta = v_2$.*

Proof: Follows from Theorem 7.1 and Theorem 6.1. $\square$

**Theorem 7.3.** *Let $R$ be a commutative ring of dimension $d$. Let us assume $R = 2R$. Let $(P\langle , \rangle)$ be a symplectic $R$-module with $P$ finitely generated projective module of even rank $\geq d$. Let $Q = (R^2 \perp P)$, and let $\widehat{Q} = (R^2 \perp Q)$. Let $\sigma \in \mathrm{Sp}(Q, \langle , \rangle)$ and $(I_2 \perp \sigma) \in \mathrm{ETrans}_{\mathrm{Sp}}(\widehat{Q}, \langle , \rangle)$. Then $\sigma$ is (stably elementary symplectic) homotopic to the identity. In fact, $\sigma = \rho(1)$, and $\rho(0) = Id$, for some $\rho(X) \in \mathrm{Sp}(Q[X], \langle , \rangle) \cap \mathrm{ETrans}_{\mathrm{Sp}}(\widehat{Q}[X], \langle , \rangle)$.*

Proof: Let us choose $\alpha(X)$ from $\mathrm{ETrans}_{\mathrm{Sp}}(\widehat{Q}[X], \langle , \rangle)$, such that $\alpha(1) = I_2 \perp \sigma$, and $\alpha(0) = Id$. Let $e_1\alpha(X) = v(X) \in \mathrm{Um}(\widehat{Q}[X], (X^2 - X)\widehat{Q}[X])$. Also $e_1 = (1, 0, 0) \in \mathrm{Um}(\widehat{Q}[X], (X^2 - X)\widehat{Q}[X])$. Therefore by Lemma 7.2, we have $\beta(X) \in \mathrm{ETrans}_{\mathrm{Sp}}(\widehat{Q}[X], (X^2-X)\widehat{Q}[X], \langle , \rangle)$, such that $v(X)\beta(X) = (1, 0, 0)$, i.e, $e_1\alpha(X)\beta(X) = e_1$. Let us call the product $\alpha(X)\beta(X) = \delta(X)$. Since $\delta(X) \in \mathrm{ETrans}_{\mathrm{Sp}}(\widehat{Q}[X], \langle , \rangle)$, we have $\langle e_1\delta(X), e_2\delta(X)\rangle = \langle e_1, e_2\rangle = 1$, and hence $e_2\ \delta(X) = (a(X), 1, q(X)) = u(X)$ (say). Note that $\delta(0) = Id$, hence $u(0) = (0, 1, 0)$. Also $\delta(1) = Id$, hence $u(1) = (0, 1, 0)$, i.e, $u(X) \equiv (0, 1, 0) \pmod{(X^2 - X)}$. Let $\mathfrak{m}$ be a maximal ideal of $R$ and $\varphi_{\mathfrak{m}}$ be the alternating matrix over $R_{\mathfrak{m}}$, which corresponds to the alternating bilinear form $\langle , \rangle$. Let us choose an element

$$\begin{aligned} \gamma(X) &= \begin{pmatrix} 1 & 0 & 0 \\ -a(X) & 1 & -q(X) \\ {\varphi_{\mathfrak{m}}^{-1}}^t q(X)^t & 0 & I \end{pmatrix} \\ &= \rho(q(X)\varphi_{\mathfrak{m}}^{-1}, -a(X)) \end{aligned}$$

from $\mathrm{ETrans}_{\mathrm{Sp}}(\widehat{Q}_{\mathfrak{m}}[X], (X^2-X)\widehat{Q}_{\mathfrak{m}}[X], \langle , \rangle)$, such that $e_1\ \gamma(X) = e_1$ and $u(X)\ \gamma(X) = e_2 = u(0)$. This is true for all maximal ideals $\mathfrak{m}$ of $R$. By Theorem 5.20 there is a $\widetilde{\gamma}(X) \in \mathrm{ETrans}_{\mathrm{Sp}}(\widehat{Q}[X], (X^2-X)\widehat{Q}[X], \langle , \rangle)$ such that $e_1\ \widetilde{\gamma}(X) = e_1$ and $u(X)\ \widetilde{\gamma}(X) = e_2$. Let us call $\delta(X)\widetilde{\gamma}(X) = \eta(X)$. Clearly $\eta(X) \in \mathrm{ETrans}_{\mathrm{Sp}}(\widehat{Q}[X], \langle , \rangle)$, and $e_1\ \eta(X) = e_1; e_2\ \eta(X) = e_2$. Let $\eta(X) = I_2 \perp \rho(X)$, where $\rho(X) = \eta(X)|_{Q[X]}$. Note that $\rho(X) \in \mathrm{Sp}(Q[X], \langle , \rangle)$, and $I_2 \perp \rho(1) = \eta(1) = \delta(1)\widetilde{\gamma}(1) = \alpha(1)\beta(1)\widetilde{\gamma}(1) = \alpha(1) = (I_2 \perp \sigma)$, and $\rho(0) = Id$. $\square$

**Theorem 7.4.** ([4], Theorem 3.13) *Let $(R, \mathfrak{m})$ be a regular local ring. Assume that $R$ contains a field, or characteristic of $R \setminus \mathfrak{m} \notin \mathfrak{m}^2$. Then*

$$\mathrm{Sp}(R[X]^{2n+2}, \langle , \rangle_{\varphi_{\mathfrak{m}}}) = \mathrm{ETrans}_{\mathrm{Sp}}(R[X]^{2n+2}, \langle , \rangle_{\varphi_{\mathfrak{m}}}),$$

*for $n \geq 1$, where $\varphi_{\mathfrak{m}}$ is the associated matrix of the alternating bilinear form $\langle , \rangle$.* $\square$

The next corollary improves Basu-Rao estimate in the module case over finitely generated rings.

**Corollary 7.5.** *Let $R$ be a finitely generated non-singular algebra of dimension $d$ over $K$, where $K$ is either a finite field or the algebraic closure of a finite field. Let us assume $R = 2R$. Let $(P.\langle , \rangle)$ be a symplectic $R$-module with $P$ finitely generated projective module of even rank $\geq d-2$. Let $Q = (R^2 \perp P)$, and $\widehat{Q} = (R^2 \perp Q)$. Let $\sigma \in \mathrm{Sp}(Q, \langle , \rangle)$ and $(I_2 \perp \sigma) \in \mathrm{ETrans}_{\mathrm{Sp}}(\widehat{Q}, \langle , \rangle)$. Then $\sigma$ belongs to* $\mathrm{ETrans}_{\mathrm{Sp}}(Q, \langle , \rangle)$.

Proof: From the proof of Theorem 7.3 it follows that $\sigma = \rho(1)$ for some $\rho(X) \in \mathrm{Sp}(Q[X], \langle , \rangle)$, with $\rho(0) = Id$. Using Theorem 7.4 we get that $\mathrm{Sp}(R_{\mathfrak{m}}[X]^{2n+2}, \langle , \rangle_{\varphi_{\mathfrak{m}}})$ $= \mathrm{ETrans}_{\mathrm{Sp}}(R_{\mathfrak{m}}[X]^{2n+2}, \langle , \rangle_{\varphi_{\mathfrak{m}}})$, for all maximal ideals $\mathfrak{m}$ of $R$. This implies $\rho(X) \in \mathrm{ETrans}_{\mathrm{Sp}}(R_{\mathfrak{m}}[X]^{2n+2}, \langle , \rangle_{\varphi_{\mathfrak{m}}})$, for all maximal ideals $\mathfrak{m}$ in $R$. By Lemma 5.19, $\rho(X) \in \mathrm{ETrans}_{\mathrm{Sp}}(R[X], \langle , \rangle)$. Hence $\sigma = \rho(1)$ belongs to $\mathrm{ETrans}_{\mathrm{Sp}}(R, \langle , \rangle)$. $\square$

# Appendix

**Alpesh Dhorajia**
Department of Mathematics, Indian Institute of Technology, Mumbai 400 076, India

The aim of this Appendix is to work out via commutator laws the size 4 cases in the study above. This entails proving Lemma 2.9 and Lemma 3.1 above in the symplectic size 4 case.

In order to prove Lemma 2.9 in the symplectic case we need to establish the following inclusion first.

**Lemma 7.6.** Let $n \geq 2$ (i.e. $2n \geq 4$), $R$ be a commutative ring with $R = 2R$, and let $I$ be an ideal of $R$. Then $\mathrm{ESp}_{2n}(R, I) \subseteq \mathrm{ESp}^1_{2n}(R, I)$.

Proof: It suffices to show $\mathrm{ESp}^1_{2n}(R, I)$ contains the set $S_{ij} = \{ge_{ij}(a)ge_{ji}(x)ge_{ij}(-a) : a \in R, x \in I\}$, for all $i, j$, with $i \neq j$. First we state the following identities

$$[gh, k] = \left({}^{g}[h,k]\right)[g,k], \tag{3}$$

$$[g, hk] = [g,h]\left({}^{h}[g,k]\right), \tag{4}$$

$${}^{g}[h,k] = [{}^{g}h, {}^{g}k], \tag{5}$$

where ${}^{g}h$ denotes $ghg^{-1}$ and $[g,h] = ghg^{-1}h^{-1}$. These identities will be used throughout the proof without mentioning it always.

Also we need the following relations which hold for all integers $i, j$ with $i \neq \sigma(j)$ and all elements $a, b \in R$.

$$\begin{aligned}
[se_{ij}(a), se_{\sigma(i)i}(b)] &= se_{\sigma(i)j}(-2ab)se_{\sigma(j)j}((-1)^{i+j}a^2b), && (6)\\
[se_{ij}(a), se_{j\sigma(j)}(b)] &= se_{i\sigma(i)}((-1)^{i+j}a^2b)se_{i\sigma(j)}(2ab), && (7)\\
[se_{\sigma(i)i}(a), se_{\sigma(j)\sigma(i)}(b)] &= se_{\sigma(j)i}(-2ab)se_{\sigma(j)j}((-1)^{i+j+1}ab^2), && (8)\\
[se_{\sigma(i)i}(a), se_{ij}(b)] &= se_{\sigma(i)j}(2ab)se_{\sigma(j)j}((-1)^{i+j+1}ab^2), && (9)\\
[se_{ij}(a), se_{j\sigma(i)}(b)] &= se_{i\sigma(i)}(2ab), && (10)\\
[se_{ij}(a), se_{\sigma(j)i}(b)] &= se_{\sigma(j)j}(-2ab), && (11)\\
[se_{\sigma(i)i}(a), se_{\sigma(j)j}(b)] &= 1, && (12)\\
[se_{\sigma(j)j}(a), se_{\sigma(i)j}(b)] &= 1, && (13)\\
se_{i\sigma(i)}(a) &= [se_{ik}(a/2), se_{k\sigma(i)}(1)], \text{where } k \neq i, \sigma(i), && (14)\\
[se_{ij}(a), se_{kl}(b)] &= 1, \text{where } i \neq \sigma(k), l \text{ and } j \neq k, \sigma(k). && (15)
\end{aligned}$$

Note that $se_{ij}(x) \in \mathrm{ESp}^1_{2n}(R, I)$, for $1 \leq i, j \leq 2$ and $x \in I$. For $3 \leq i, j \leq 2n$ we have $g_{ij}(x) = [ge_{i1}(x/2), ge_{1j}(1)] \in \mathrm{ESp}^1_{2n}(R, I)$. In the following computation we will express generators of $S_{ij}$ in terms of $g_{ij}(x)$ or $ge_{1j}(a)$, where $x \in I, a \in R$. Also note that we will use $*$ to represent an element of the ideal $I$ and $\circledast$ to represent an element of the ring $R$, which does not belong to the ideal $I$.

*Case (a):* Let us assume $i \neq \sigma(j)$ and $1 \leq i \leq 2$, $3 \leq j \leq 2n$. If $i = 1$ it is trivial. We assume $i = 2$. Using the equations above we get

$$\begin{aligned}
{}^{se_{ij}(a)}se_{ji}(x) &= {}^{se_{ij}(a)}\big([se_{j\sigma(i)}(*), se_{\sigma(i)i}(1)]se_{j\sigma(j)}(*)\big)\\
&= [{}^{se_{ij}(a)}se_{j\sigma(i)}(*), {}^{se_{ij}(a)}se_{\sigma(i)i}(1)]\,{}^{se_{ij}(a)}se_{j\sigma(j)}(*)\\
&= [se_{i\sigma(i)}(*)se_{j\sigma(i)}(*), se_{\sigma(i)i}(1)se_{\sigma(i)j}(\circledast)se_{\sigma(j)j}(\circledast)]\,{}^{se_{ij}(\circledast)}se_{j\sigma(j)}(*)\\
&= [se_{i\sigma(i)}(*)se_{j\sigma(i)}(*), se_{\sigma(j)j}(\circledast)]\ \big({}^{se_{\sigma(j)j}(\circledast)}[se_{i\sigma(i)}(*)se_{j\sigma(i)}(*),\\
&\quad se_{\sigma(i)i}(1)se_{\sigma(i)j}(\circledast)]\big)\ {}^{se_{ij}(\circledast)}se_{j\sigma(j)}(*)\\
&= ABC\ (say),
\end{aligned}$$

where

$$\begin{aligned}
A &= [se_{i\sigma(i)}(*)se_{j\sigma(i)}(*), se_{\sigma(j)j}(\circledast)],\\
B &= {}^{se_{\sigma(j)j}(\circledast)}[se_{i\sigma(i)}(*)se_{j\sigma(i)}(*), se_{\sigma(i)i}(1)se_{\sigma(i)j}(\circledast)],\\
C &= {}^{se_{ij}(a)}se_{j\sigma(j)}(*).
\end{aligned}$$

Now

$$\begin{aligned}
A &= [se_{i\sigma(i)}(*)se_{j\sigma(i)}(*), se_{\sigma(j)j}(\circledast)]\\
&= se_{i\sigma(i)}(*)se_{j\sigma(i)}(*)se_{\sigma(j)j}(\circledast)se_{j\sigma(i)}(*)^{-1}se_{i\sigma(i)}(*)^{-1}se_{\sigma(j)j}(\circledast)^{-1}\\
&= se_{i\sigma(i)}(*)se_{j\sigma(i)}(*)^{se_{\sigma(j)j}(\circledast)}se_{j\sigma(i)}(*)^{se_{\sigma(j)j}(\circledast)}se_{i\sigma(i)}(*)\\
&= se_{i\sigma(i)}(*)se_{j\sigma(i)}(*)\big(se_{j\sigma(i)}(*)se_{\sigma(j)\sigma(i)}(*)se_{i\sigma(i)}(*)\big)se_{i\sigma(i)}(*),
\end{aligned}$$

$$\begin{array}{rcl} B & = & {}^{se_{\sigma(j)j}(\circledast)}[se_{i\sigma(i)}(*)se_{j\sigma(i)}(*), se_{\sigma(i)i}(1)se_{\sigma(i)j}(\circledast)] \\ & = & [{}^{se_{\sigma(j)j}(\circledast)}se_{i\sigma(i)}(*){}^{se_{\sigma(j)j}(\circledast)}se_{j\sigma(i)}(*), {}^{se_{\sigma(j)j}(\circledast)}\, se_{\sigma(i)i}(1){}^{se_{\sigma(j)j}(\circledast)}se_{\sigma(i)j}(\circledast)] \\ & = & [se_{i\sigma(i)}(*)se_{j\sigma(i)}(*)se_{i\sigma(i)}(*)se_{\sigma(j)\sigma(i)}(*), se_{\sigma(i)i}(1)se_{\sigma(i)j}(\circledast)], \\ C & = & se_{j\sigma(j)}(*)se_{i\sigma(i)}(*)se_{i\sigma(j)}(*). \end{array}$$

It is clear that $A, C \in \mathrm{ESp}^1_{2n}(R,I)$. In this case $\sigma(i) = 1$, and hence $B \in \mathrm{ESp}^1_{2n}(R,I)$.

*Case (b):* Let us assume $i \neq \sigma(j)$ and $3 \leq i, j \leq 2n$. From the calculation in Case (a) it is clear that $A, C \in \mathrm{ESp}^1_{2n}(R,I)$. We need to show $B \in \mathrm{ESp}^1_{2n}(R,I)$ in this case. We have

$$\begin{array}{rcl} B & = & [se_{i\sigma(i)}(*)se_{j\sigma(i)}(*)se_{i\sigma(i)}(*)se_{\sigma(j)\sigma(i)}(*), se_{\sigma(i)i}(1)se_{\sigma(i)j}(\circledast)] \\ & = & [se_{j\sigma(i)}(*)se_{i\sigma(i)}(*)se_{\sigma(j)\sigma(i)}(*), se_{\sigma(i)i}(1)se_{\sigma(i)j}(\circledast)] \\ & = & se_{j\sigma(i)}(*)se_{i\sigma(i)}(*)se_{\sigma(j)\sigma(i)}(*){}^{se_{\sigma(i)i}(1)}\big({}^{se_{\sigma(i)j}(\circledast)}se_{\sigma(j)\sigma(i)}(*)\ {}^{se_{\sigma(i)j}(\circledast)}se_{i\sigma(i)}(*) \\ & & {}^{se_{\sigma(i)j}(\circledast)}se_{j\sigma(i)}(*)\big) \\ & = & se_{j\sigma(i)}(*)se_{i\sigma(i)}(*)se_{\sigma(j)\sigma(i)}(*){}^{se_{\sigma(i)i}(1)}\big(se_{\sigma(j)\sigma(i)}(*)se_{\sigma(j)j}(*)\big) \\ & & {}^{se_{\sigma(i)i}(1)}\big(se_{ij}(*)se_{i\sigma(i)}(*)se_{\sigma(j)j}(*)\big) \\ & & {}^{se_{\sigma(i)i}(1)}\big(se_{j1}(*)se_{\sigma(i)1}(*)se_{1\sigma(i)}(1)se_{1j}(a)se_{\sigma(i)1}(*)se_{j1}(*)se_{1j}(\circledast)se_{1\sigma(i)}(-1)\big) \end{array}$$

From the computation below it will be clear that $B \in \mathrm{ESp}^1_{2n}(R,I)$. We execute the calculation with the help of equations (8), (9), (12) and (15).

$$\begin{array}{rcl} {}^{se_{\sigma(i)i}(1)}se_{\sigma(j)\sigma(i)}(*) & = & se_{\sigma(j)\sigma(i)}(*)se_{\sigma(j)i}(-*)se_{\sigma(j)j}(*) \\ {}^{se_{\sigma(i)i}(1)}se_{\sigma(j)j}(*) & = & se_{\sigma(j)j}(*) \\ {}^{se_{\sigma(i)i}(1)}se_{ij}(*) & = & se_{ij}(*)se_{\sigma(i)j}(a*)se_{\sigma(j)j}(*) \\ {}^{se_{\sigma(i)i}(1)}se_{i\sigma(i)}(*) & = & {}^{se_{\sigma(i)i}(1)}[se_{i1}(*), se_{1\sigma(i)}(1)] \\ & = & [se_{i1}(*)se_{\sigma(i)1}(*)se_{21}(*), se_{1\sigma(i)}(1)se_{1i}(-1)se_{12}(1')] \\ {}^{se_{\sigma(i)i}(1)}se_{1j}(\circledast) & = & se_{1j}(\circledast) \\ {}^{se_{\sigma(i)i}(1)}se_{j1}(*) & = & se_{j1}(*) \\ {}^{se_{\sigma(i)i}(1)}se_{\sigma(i)1}(*) & = & se_{\sigma(i)1}(*) \\ {}^{se_{\sigma(i)i}(1)}se_{1\sigma(i)}(1) & = & se_{1\sigma(i)}(1)se_{1i}(-1)se_{12}(-1). \end{array}$$

*Case (c):* Let us assume $i = \sigma(j)$. Clearly $S_{12} \subseteq \mathrm{ESp}^1_{2n}(R,I)$. When $i \geq 3$ we have

$$\begin{array}{rcl} {}^{se_{ij}(a)}se_{ji}(x) & = & {}^{se_{ij}(a)}[se_{j1}(x/2), se_{1i}(1)] \\ & = & [se_{j1}(*)se_{i1}(*)se_{21}(*), se_{1i}(\circledast)se_{1j}(\circledast)se_{12}(\circledast), \end{array}$$

and hence $S_{i\sigma(i)} \subseteq \mathrm{ESp}^1_{2n}(R,I)$, for $i \geq 3$.

Now we consider the case $i = \sigma(j)$ and $i = 2$. For an integer $k \neq 1, 2$ we have

$$\begin{aligned}
{}^{se_{21}(a)}se_{12}(x) &= se_{21}(a)se_{12}(x)se_{21}(-a) \\
&= {}^{se_{21}(a)}[se_{1k}(1), se_{k2}(*)] \\
&= [{}^{se_{21}(a)}se_{1k}(1), {}^{se_{21}(a)}se_{k2}(*)] \\
&= [\alpha, se_{k1}(*)se_{k\sigma(k)}(*)se_{k2}(*)] \\
&= {}^{\alpha}se_{k1}(*)\,{}^{\alpha}se_{k\sigma(k)}(*)\,{}^{\alpha}se_{k2}(*)se_{k1}(*)se_{k\sigma(k)}(*)se_{k2}(*),
\end{aligned}$$

where $\alpha = {}^{se_{21}(a)}se_{1k}(1)$ which can be expressed in the following three different ways, namely

$$\begin{aligned}
\alpha &= se_{1k}(1)se_{2k}(\circledast)se_{\sigma(k)k}(\circledast) \\
&= se_{\sigma(k)k}(\circledast)se_{2k}(\circledast)se_{1k}(1) \\
&= se_{2k}(\circledast)se_{\sigma(k)k}(\circledast)se_{1k}(1),
\end{aligned}$$

and we will use them as we find it convenient. Now

$$\begin{aligned}
{}^{\alpha}se_{k1}(*) &= {}^{se_{1k}(1)se_{2k}(\circledast)se_{\sigma(k)k}(\circledast)}se_{k1}(*) \\
&= {}^{se_{1k}(1)se_{2k}(\circledast)}\big(se_{\sigma(k)k}(\circledast)se_{k1}(*)se_{\sigma(k)k}(\circledast)^{-1}\big) \\
&= {}^{se_{1k}(1)se_{2k}(\circledast)}se_{k1}(*) \\
&= {}^{se_{1k}(1)}\big(se_{2k}(\circledast)se_{k1}(*)se_{2k}(\circledast)^{-1}\big) \\
&= {}^{se_{1k}(1)}\big(se_{21}(*)se_{k1}(*)\big) \\
&= se_{2k}(*)se_{\sigma(k)k}(*)se_{21}(*)se_{1k}(1)se_{k1}(*)se_{1k}(1)^{-1}.
\end{aligned}$$

Given that $*$ represents elements of $I$ and $S_{1k} \subseteq \mathrm{ESp}^1_{2n}(R, I)$, we have ${}^{\alpha}se_{k1}(*) \in \mathrm{ESp}^1_{2n}(R, I)$.

$$\begin{aligned}
{}^{\alpha}se_{k\sigma(k)}(*) &= {}^{se_{\sigma(k)k}(\circledast)se_{2k}(\circledast)se_{1k}(1)}se_{k\sigma(k)}(*) \\
&= {}^{se_{\sigma(k)k}(\circledast)se_{2k}(\circledast)}\big(se_{1k}(1)se_{k\sigma(k)}(*)se_{1k}(1)^{-1}\big) \\
&= {}^{se_{\sigma(k)k}(\circledast)se_{2k}(\circledast)}\big(se_{12}(*)se_{k2}(*)se_{k\sigma(k)}(*)\big) \\
&= {}^{se_{\sigma(k)k}(\circledast)se_{2k}(\circledast)}se_{12}(*)\ {}^{se_{\sigma(k)k}(\circledast)se_{2k}(\circledast)}se_{k2}(*)\ {}^{se_{\sigma(k)k}(\circledast)se_{2k}(\circledast)}se_{k\sigma(k)}(*) \\
&= {}^{se_{\sigma(k)k}(\circledast)}\big(se_{1k}(*)se_{\sigma(k)k}(*)se_{12}(*)\big)\ {}^{se_{2k}(\circledast)se_{\sigma(k)k}(\circledast)}se_{k2}(*) \\
&\quad\ {}^{se_{\sigma(k)k}(\circledast)}\big(se_{21}(*)se_{k1}(*)se_{k\sigma(k)}(*)\big) \\
&= \big(se_{1k}(*)se_{\sigma(k)k}(*)se_{12}(*)\big)\ {}^{se_{2k}(\circledast)}\big(se_{1k}(*)se_{12}(*)se_{k2}(*)\big) \\
&\quad \big(se_{21}(*)se_{2k}(*)se_{21}(*)se_{k1}(*)se_{\sigma(k)k}(\circledast)se_{k\sigma(k)}(*)se_{\sigma(k)k}(\circledast)^{-1}\big) \\
&= \big(se_{1k}(*)se_{\sigma(k)k}(*)se_{12}(*)\big) \\
&\quad \big(se_{\sigma(k)k}(*)se_{1k}(*)se_{\sigma(k)k}(*)se_{12}(*)se_{2k}(\circledast)se_{k2}(*)se_{2k}(\circledast)^{-1}\big) \\
&\quad \big(se_{21}(*)se_{2k}(*)se_{21}(*)se_{k1}(*)se_{\sigma(k)k}(\circledast)se_{k\sigma(k)}(*)se_{\sigma(k)k}(\circledast)^{-1}\big)
\end{aligned}$$

Given that $*$ represents elements of $I$ and $S_{2k}, S_{\sigma(k)k} \subseteq \mathrm{ESp}^1_{2n}(R,I)$, we have ${}^{\alpha}se_{k\sigma(k)}(*) \in \mathrm{ESp}^1_{2n}(R,I)$.

$$\begin{array}{rcl}
{}^{\alpha}se_{k2}(*) & = & {}^{se_{2k}(\circledast)se_{\sigma(k)k}(\circledast)se_{1k}(1)}se_{k2}(*) \\
& = & {}^{se_{2k}(\circledast)se_{\sigma(k)k}(\circledast)}\big(se_{12}(*)se_{k2}(*)\big) \\
& = & {}^{se_{2k}(\circledast)}\big(se_{12}(*)se_{1k}(*)se_{21}(*)se_{k2}(*)\big) \\
& = & se_{1k}(*)se_{\sigma(k)k}(*)se_{12}(*)se_{\sigma(k)k}(*)se_{1k}(*)se_{21}(*)se_{2k}(\circledast) \\
& & se_{k2}(*)se_{2k}(\circledast)^{-1}
\end{array}$$

Similarly as above $*$ represents elements of $I$ and $S_{2k} \subseteq \mathrm{ESp}^1_{2n}(R,I)$ implies ${}^{\alpha}se_{k2}(*)$ is in $\mathrm{ESp}^1_{2n}(R,I)$. Therefore $S_{21} \subseteq \mathrm{ESp}^1_{2n}(R,I)$. $\square$

Proof of Lemma 2.9: Let $f : \mathrm{E}^1(n,R,I) \longrightarrow \mathrm{E}^1(n,R/I,0)$. Note that

$$\ker(f) \quad \subseteq \quad \mathrm{E}^1(n,R,I) \cap \mathrm{G}(n,R,I).$$

Let $M = \prod ge_{j1}(x_j)ge_{1i}(a_i) \in \mathrm{E}^1(n,R,I) \cap \mathrm{G}(n,R,I)$. Note that $M \in \mathrm{G}(n,R,I)$ implies $\bar{M} = \prod ge_{j1}(0)ge_{1i}(\bar{a_i}) = \prod ge_{1i}(\bar{a_i}) = I_n$, i.e, $M \in \ker(f)$. Therefore $\ker(f) = \mathrm{E}^1(n,R,I) \cap \mathrm{G}(n,R,I)$.

Now we shall prove that $\ker(f) = \mathrm{E}(n,R,I)$. Let $E = \prod_{k=1}^{r} ge_{j_k1}(x_k)ge_{1i_k}(a_k) \in \ker(f)$. Note that $E$ can be written as $ge_{j_11}(x_1)\prod_{k=2}^{r}\gamma_k ge_{j_k1}(x_k)\gamma_k^{-1}$, where $\gamma_l$ is equal to $\prod_{k=1}^{l-1}\ ge_{1i_k}(a_k) \in \mathrm{E}(n,R)$, and hence $\ker(f) \subseteq \mathrm{E}(n,R,I)$. The reverse inclusion follows from the fact that $\mathrm{E}(n,R,I) \subseteq \mathrm{E}^1(n,R,I)$. $\square$

Proof of Lemma 3.1 in the symplectic case when $n = 4$: Given that $se_{ij}(Xf(X))$ is in $\mathrm{ESp}^1_{2n}(R[X], I[X])$. First assume $i = 1$ and $f(X) \in R[X]$. We prove the result using induction on $r$, the number of generators of $\varepsilon$. Let $r = 1$ and $\varepsilon = se_{pq}(a)$. Note that when $p = 1, a \in R$, and when $q = 1, a \in I$. Also note that we will use $*$ to represent an element of the ideal $I$.

*Case (1):* Let $(p,q) = (1,j))$. In this case

$$se_{1j}(a)\ ge_{1j}(Y^2Xf(Y^2X))\ se_{1j}(-a) \quad = \quad se_{1j}(Y^2Xf(Y^2X)).$$

*Case (2):* Let $(p,q) = (1,\sigma(j))$. In this case

$$\begin{array}{rl}
& se_{1\sigma(j)}(a)\ se_{1j}(Y^2Xf(Y^2X))\ se_{1\sigma(j)}(-a) \\
= & se_{12}(*Y^2Xf(Y^2X))se_{1j}(Y^2Xf(Y^2X)).
\end{array}$$

*Case (3):* Let $(p,q) = (1,k)$ where $k \neq j, \sigma(j)$. In this case

$$se_{1k}(a)\ se_{1j}(Y^2Xf(Y^2X))\ se_{1k}(-a) \quad = \quad se_{1j}(Y^2Xf(Y^2X)).$$

*Case (4):* Let $(p,q) = (2,1)$ and $j = 2$. Choose $k \neq 1, 2$. Then

$$\begin{array}{rl}
& se_{21}(a)\ se_{12}(Y^2Xf(Y^2X))\ se_{21}(-a) \\
= & se_{\sigma(k)1}(*Y)\ se_{k1}(*Y).
\end{array}$$

*Case (5):* Let $(p,q)=(2,1)$ and $j\neq 2$. In this case

$$\begin{aligned} & se_{21}(a)\ se_{1j}(Y^2Xf(Y^2X))\ se_{21}(-a) \\ = \ & se_{\sigma(j)1}(*Y)\ [se_{\sigma(j)1}(*Y), se_{1j}(Y^3X^2f(Y^2X))]. \end{aligned}$$

*Case (6):* Let $(p,q)=(k,1)$, where $k\neq 2$ and $j=2$. In this case

$$\begin{aligned} & se_{k1}(a)\ se_{12}(Y^2Xf(Y^2X))\ se_{k1}(-a) \\ = \ & se_{1\sigma(k)}(*Y^2Xf(Y^2X))\ se_{k1}(*Y)\ se_{1\sigma(k)}(Y^3Xf(Y^2X)) \\ & se_{12}(Y^2Xf(Y^2X)). \end{aligned}$$

*Case(7):* Let $(p,q)=(\sigma(j),1)$ and $j\neq 2$. In this case

$$\begin{aligned} & se_{\sigma(j)1}(a)\ se_{1j}(Y^2Xf(Y^2X))\ se_{\sigma(j)1}(-a) \\ = \ & [se_{\sigma(j)1}(*Y), se_{1j}(YXf(Y^2X))]\ se_{1j}(Y^2Xf(Y^2X)). \end{aligned}$$

*Case(8):* Let $(p,q)=(j,1)$ and $j\neq 2$. In this case

$$\begin{aligned} & se_{j1}(a)\ se_{1j}(Y^2Xf(Y^2X))\ se_{j1}(-a) \\ = \ & se_{j1}(a)\ se_{1j}(Y^2Xf(Y^2X))\ se_{12}((-1)^{j+2}Y^3Xf(Y^2X)/2) \\ & \ se_{12}((-1)^{j+1}Y^3Xf(Y^2X)/2)\ se_{j1}(-a) \\ = \ & se_{j1}(a)\ [se_{\sigma(j)j}(-xyf(Y^2X)/2), se_{\sigma(j)1}(Y)] \\ & \ se_{12}((-1)^{j+1}Y^3Xf(Y^2X)/2)\ se_{j1}(-a) \\ = \ & [se_{\sigma(j)1}(aYXf(Y^2X))\ se_{21}((-1)^j a^2YXf(Y^2X))\ se_{\sigma(j)j}(-YXf(Y^2X)/2), \\ & se_{21}(2aY)\ se_{\sigma(j)1}(Y)]\ \ se_{\sigma(j)j}(a^2Y^3Xf(Y^2X)/2) \\ & se_{j2}((-1)^{j+1}(aY^3Xf(Y^2X))\ se_{12}((-1)^{j+1}Y^3Xf(Y^2X)/2) \\ = \ & se_{\sigma(j)1}(*YXf(Y^2X))\ se_{21}(*YXf(Y^2X))\ se_{21}(*Y^2)\ se_{21}(*YXf(Y^2X)/2) \\ & se_{\sigma(j)1}(*YXf(Y^2X))\ [se_{\sigma(j)1}(*Y), se_{1j}(Y^2Xf(Y^2X)/2)] \\ & se_{1\sigma(j)}(*Y^3Xf(Y^2X))\ se_{12}((-1)^{j+1}Y^3Xf(Y^2X)/2). \end{aligned}$$

Arguing as in the proof of Lemma 3.1 of §3, we can complete this proof. □

Using equations (6) to (15) we can improve Theorem 5.6 of [5]. For this purpose we first establish the following lemma ([5], Lemma 5.2) for size 4:

**Lemma 7.7.** *Let $I$ be an ideal of $R$. Assume that $2R=R$. Then $\mathrm{ESp}_{2n}(R,I^2)\subset \mathrm{ESp}_{2n}(I)$, for $n\geq 2$.*

Proof: Let $z^*=\sum a_tb_t$ with $a_t,b_t\in I$. Let $\beta=se_{ij}(z^*)\in \mathrm{ESp}_{2n}(I^2)$ and $\alpha=se_{kl}(z)\in \mathrm{ESp}_{2n}(R)$ for some $z\in R$. It suffices to show that $\alpha\beta\alpha^{-1}\in \mathrm{ESp}_{2n}(I)$. If $(k,l)\neq(j,i)$ and $(k,l)\neq(\sigma(i),\sigma(j))$, then the matrix $\alpha\beta\alpha^{-1}$ splits into a product of elementary matrices from $\mathrm{ESp}_{2n}(I)$.

When $(k,l) = (j,i)$ or $(k,l) = (\sigma(i),\sigma(j))$, we have

$$\begin{aligned}
\alpha\beta\alpha^{-1} &= \alpha\ \big(\prod_t se_{ij}(a_tb_t)\big)\ \alpha^{-1} \\
&= \alpha\ \big(\prod_t se_{ij}(a_tb_t)\ se_{\sigma(j)j}((-1)^{i+j}a_tb_t^2/2)\ se_{\sigma(j)j}((-1)^{i+j+1}a_tb_t^2/2)\big)\ \alpha^{-1} \\
&= \alpha\ \big(\prod_t [se_{\sigma(i)j}(b_t),\ se_{i\sigma(i)}(-a_t/2)]\ se_{\sigma(j)j}((-1)^{i+j+1}a_tb_t^2/2)\big)\ \alpha^{-1} \\
&= \prod_t [se_{\sigma(i)i}(-2b_tz)\ se_{\sigma(i)j}(b_t),\ se_{j\sigma(j)}((-1)^{i+j+1}a_tz^2/2)\ se_{j\sigma(i)}(-a_tz) \\
&\quad se_{i\sigma(i)}(-a_t/2)]\ se_{\sigma(j)i}((-1)^{i+j}a_tb_t^2z)\ se_{\sigma(i)i}(-a_tb_t^2z^2/2) \\
&\quad se_{\sigma(j)j}((-1)^{i+j+1}a_tb_t^2/2),
\end{aligned}$$

and hence $\alpha\beta\alpha^{-1} \in \mathrm{ESp}_{2n}(I)$. ☐

Now we have Lemma 5.1 to Lemma 5.4 and Theorem 5.5 of [5] for Elementary symplectic group of size 4, which will prove the size 4 case of ([5], Theorem 5.6), *viz.*:

**Theorem 7.8.** ([5], Theorem 5.6) *Let $I$ be an ideal of $R$. Assume $2R = R$. Then the natural map $\mathrm{Um}_{2n}(R,I)/\mathrm{ESp}_{2n}(R,I) \longrightarrow \mathrm{Um}_{2n}(R,I)/\mathrm{E}_{2n}(R,I)$ is bijective for $n \geq 2$.* ☐

# References

[1] H. Bass, $K$-theory and stable algebra, Inst. Publ. Math. IHES 22 (1964) 5–60.

[2] H. Bass, Algebraic $K$-theory, W.A. Benjamin, Inc, 1968.

[3] H. Bass, Unitary algebraic $K$-theory, Lecture Notes in Mathematics 343 (1973) 57–265

[4] R. Basu, R.A. Rao, Injective Stability for $K_1$ of Classical Modules, Journal of Algebra 323 no. 4 (2010) 867–877.

[5] P. Chattopadhyay, R.A. Rao, Elementary symplectic orbits and improved $K_1$-stability, Journal of $K$-Theory 7 (2011) 389–403.

[6] Hazrat, Roozbeh; Vavilov, Nikolai $K_1$ of Chevalley groups are nilpotent. J. Pure Appl. Algebra 179 (2003), no. 1-2, 99–116.

[7] Hazrat, R.; Petrov, V.; Vavilov, N. Relative subgroups in Chevalley groups. J. K-Theory 5 (2010), no. 3, 603–618.

[8] W. van der Kallen, A group structure on certain orbit sets of unimodular rows, Journal of Algebra 82 no. 2 (1983) 363–397.

[9] S. Mandal, Projective module and complete intersection, Lecture Notes in Mathematics 1672 (1997).

[10] D. Quillen, Projective modules over polynomial rings, Invent. Math. 36 (1976) 167–171.

[11] R.A. Rao, An elementary transformation of a special unimodular vector to its top coefficient vector, Proceedings of the American Mathematical Society 93 no. 1 (1985) 21–24.

[12] A.A. Suslin, L.N. Vaserstein, Serre's problem on Projective Modules over Polynomial Rings and Algebraic K-theory, Math. USSR Izvestija 10 (1976) 937–1001.

[13] A.A. Suslin, On the Structure of the Special Linear Group over Polynomial Rings, Math. USSR. Izvestija 11 (1977) 221–238.

[14] R.G. Swan, Serre's Problem, Queen's paper in pure and applied mathematics 42 Queen's University, Kingston, Ontario (1975) 1–60.

[15] L.N. Vaserstein, On the normal subgroups of $\mathrm{GL}_n$ over a ring, Algebraic $K$-theory, Evanston 1980 (Proc. Conf., Northwestern Univ., Evanston, Ill., 1980) 456–465; Lecture Notes in Math. 854, Springer, Berlin-New York, 1981.